

Variety and Seed Study and Control Group







| GEVES in a few words | 3 |
|---|----|
| Role and quality | 4 |
| Place an order | 5 |
| Your contacts at GEVES | 6 |
| Supply of samples to the SNES | 7 |
| Order an analysis | 9 |
| All Species | 10 |
| Seed mixture species | 14 |
| Vegetables | 15 |
| Ornamental and Fruit crops | 31 |
| Aromatic medicinal | 33 |
| Micro-cleaning | 35 |
| Cytology | 36 |
| Radiography 2D and tomography | 37 |
| Biostimulation, Biocontrol, evaluation of treatment | 38 |
| Disease test supplies : inoculum and reference material | 39 |
| Sector support | 40 |
| Terms and Conditions | 41 |
| Publications | 42 |

GEVES: A unique &

GEVES is a Public Interest Group with three founding partner organisations:



• The French National Research Institute for Agriculture, Food and Environment (INRAE)- 60%



• The French Ministry of Agriculture and Food (MAA) - 20%



The French Interprofessional Organisation for Seeds and Plants - 20%

This unique set-up ensures GEVES's **independence** and **neutrality** in carrying out its activities in accordance with its regulatory and official missions and mandates. The union of state, research and sector expertise ensures that all aspects of the sector are fully taken into account.

Governance of GEVES

GEVES's Executive Board of Directors is composed of 13 members:

- 6 representatives from INRAE
- 2 representatives from the Ministry of Agriculture and Food
- 2 representatives from GNIS
- ${\it 2 staff representatives from GEVES}\\$
- The President of the CTPS

as well as a government controller (Ministry of Research) and a State Controller.

Organisation of GEVES's operating divisions

4 labs

SEV Variety Studies Department Station

Seed Testing Station

L'Anjouère O Brion Angers-Beaucouzé

L'Anjouère O Angers-Beaucouzé

L'Anjouère O Clermont-Ferrand

L'Anjouère O Cavaillon Carpentras

3 Units

O Cavaillon Carpentras

O Cavaillon Carpentras

GEVES's missions

GEVES has official, regulatory missions and carries out testing activities and methodological development which is necessary for:

- National listing of new varieties in the Official French Catalogue
- ▶ Plant variety protection
- ▶ Official seed testing as part of its NRL mandates for seeds, GMOs. and plant health (RNQPmatrix seeds)

GEVES is also responsible for the national coordination of plant genetic resources on behalf of the Ministry of Agriculture.

GEVES is the National Reference Laboratory for:

- ▶ GMO detection: GMOs in maize (seed) and soya, rapeseed and flax (seed and vegetative parts) by Decree of 19 octobre 2015
- > quality testing of seeds and propagating material by Decree of 1 March 2017
- ▶ in the field of plant health by Decree of 20 November 2020

GEVES is an approved laboratory for certain seed health quality tests

GEVES is accredited by ISTA for all species. It carries out official testing, particularly for seed exports: for phytosanitary passports and certificates as well as Orange and Blue International Certificates (OIC and BIC).

GEVES makes its specialised expertise openly available to the plant and seed sectors, providing high-quality services to a range of private customers.

Activities

To carry out its missions, GEVES performs a wide range of activities:

- Description of varieties and evaluation of genetic progress
- ▶ Quality testing for seeds and seedlings
- ▶ Methodological research
- Management of plant genetic resources
- ▶ Training courses
- ▶ Exams
- ► Consulting and expertise
- ▶ International cooperation
- ▶ Monitoring of the French network of seed testing laboratories
- ▶ Organisation of Proficiency Tests (PT)
- ► Communication
- ▶ Expertise
- ▶ Inoculum production
- Analysis to evaluate the efficiency of treatment products





Quality, Recognition & Accreditation

GEVES benefits from a global and harmonised Quality Management System.

▶ Certification ISO 9001: version 2015 - BioGEVES and VCUS variety testing (Value for Cultivation, Use

▶ Accreditation of GEVES's SNES and BioGEVES laboratories by COFRAC according to ISO 17025

GEVES is recognised as follows:

and Sustainability) since 2009





- GEVES Beaucouzé: COFRAC N°1-1316 (since 2002).
 - GEVES Le Magneraud: COFRAC N°1-6176 (since 2004).
- ▶ Accreditation by ISTA since 2001 (N°FRDL0200) for seed testing
- ▶ Entrusted by the CPVO for DUS variety testing since 2012.



Seed quality testing **SNES**



ORDER YOUR ANALYSE ONLINE

http://dsn.geves.info

- Enter your order
- Print the order summary and attach it to to your sample

For faster processing of your request, please order online



SEND YOUR ORDER VIA POST

- Complete the form corresponding to your order (OIC request or analysis order form) and join the form to your sample
- Send the sample to:

GEVES - Service clients SNES 3 rue Henri Becquerel - CS 90024 49071 Beaucouzé Cedex FRANCE

Biomolecular and biochemical testing

BioGEVES



ORDER YOUR ANALYSE ONLINE

biogeves.analyses@geves.fr



SEND YOUR ORDER VIA POST

• Send the sample to:

Detection Unit

BioGEVES

3 rue Henri Becquerel - CS 90024 49071 Beaucouzé Cedex FRANCE

Genotyping/Biochemistry Unit

BioGEVES - Le Magneraud

CS 40052 - Saint-Pierre d'Amilly 17 700 Surgères FRANCE

Variety testing at the **SEV**



REQUEST A DENOMINATION TEST



REQUEST A FIELD TEST DUS (Distinction Uniformity Stability)

christelle.godin@geves.fr

celine.delarue@geves.fr

GEVES - Service clients SEV 25 rue Georges Morel - CS 90024 49071 Beaucouzé Cedex FRANCE

Your contacts at GEVES

To contact a GEVES staff member by email: firstname.surname@geves.fr - area code number: +33(0).

Sector support Training courses, ILC, Audits

SNES Management



SNES Director Clotilde Polderman-Roussille +33 (0)2 41 22 58 10



Assistant Estelle Bertel +33 (0)2 41 22 58 02

SNES Customer Services

service.clients@geves.fr



Head of Customer Service and Sampling Alice Richard Jolly: +33 (0)2 41 22 58 13

Customer Service SNES

+33 (0)2 41 22 58 28

Thibaut Decourcelle +33 (0)2 41 22 58 17



Fabienne Brun +33 (0)2 41 22 58 91

Inr.semences@geves.fr

eil.semences@geves.fr

TRAINING COURSES : formation.semences@

ES/

ILC:

geves.fr

AUDITS:



Head of Physical Analysis Laboratory Aurélie Charrier: +33 (0)2 41 22 58 40

Radiography 2D/3D
 Purity, micro-cleaning
 Water content
 Botanic
 Sherif Hamdy +33 (0)2 41 22 58 30
 Philippe Pannetier +33 (0)2 41 22 58 43
 Céline Herbert +33 (0)2 41 22 58 30
 Diogo Tobolski +33 (0)2 41 22 58 94



Head of Germination Laboratory Sylvie Ducournau: +33 (0)2 41 22 58 70

Floral, vegetable, woody, pulses and Valérie Blouin +33 (0)2 41 22 58 78 forest species

Beetroot, vegetable, forage grasses
 Pierre Soufflet
 +33 (0)2 41 22 58 82

Agricultural crop species
 Philippe Garreau +33 (0)2 41 22 58 77



Head of Pathology Laboratory Jaiana Malabarba: +33 (0)2 41 22 58 50

Seed health Isabelle Serandat Laurent Guyot +33 (0)2 41 22 58 54 (0)2 41 22 58 59
 Variety resistance Sophie Perrot +33 (0)2 41 22 58 58
 Seed treatment evaluation Geoffrey Orgeur +33 (0)2 41 22 58 56

audit.semences@ geves.fr

Caroline

Le Quilliec

+33 (0)5 17 06 96 12

BioGEVES:

biogeves.analyses@geves.fr



Head of BioGEVES René Mathis +33 (0)2 41 22 58 34



Customer relationship Caroline Le Quilliec +33 (0)5 17 06 96 12



Contacts SEV:

Detection Unit Amandine Lê Van +33 (0)2 41 22 58 39



Biochemistry Unit Patricia Lem +33 (0)5 17 06 96 13



Genotyping Unit Arnaud Remay +33 (0)5 17 06 96 17

Rachel Tessier +33 (0)2 41 22 85 93



Head of SEV Fabien Masson +33 (0)2 41 22 85 91

SEV Customer Service



Céline Delarue +33 (0)2 41 22 86 00 (field trials)



Christelle GODIN +33 (0)2 41 22 86 93 (Denomination tests)

BioGEVES

Supply of samples to the SNES



The following information, listed on the SNES order form, is essential for processing seed samples:

- Treated seed and trade name of product. No treated sample will be accepted for analysis without this information.
- Thousand Seed Weight (TSW). This information is necessary to calculate the weight of sub-samples for bacteriology, mycology and virology. If this information is not indicated, it will be invoiced.
- Sample size. Unless indicated differently, the sample size to be provided is expressed in number of seeds.

The sample size indicated is the minimum size set by the method (larger sizes can be offered).

If you do not have the quantity requested and wish to have the analysis done on all the seeds sent, you must indicate this in your request.

Otherwise, the analysis will be put on hold, and we will contact you. You can then:

- send a new sample
- give us your agreement to carry out the analysis on all the seeds supplied.

Please take care to send your seeds in anonymous boxes and/or paper sachets without any labels or commercial names.

The analyses are not performed on GMO samples.

If you wish to make an analysis that is not listed in the price list (species, particular methods, etc.), contact Customer Services who will define with you the work that is adapted to your needs, the feasibility and the cost.



The SNES always works in compliance with the ISTA Rules, offering the same level of reliability of results, whatever the final certificate requested.

Physical quality: provide the minimum weights prescribed by the ISTA Rules, chapter 2.5.4.5. If a counting analysis is requested, provide the weight listed in table 2C column 3. If more than one counting analysis is requested on the same submitted sample, provide the quantities required to perform all the countings.

If only a purity test is requested, provide the seed quantities for the submitted sample according to the following table:

| Weight of working sample for purity analysis alone (Table 2C column 4) | Minimum weight of submitted sample for purity analysis (Table Column 4) |
|--|---|
| Between 500g and 1000g | Minimum working sample weight for purity analysis + 100g |
| Under 500g | 2,5 times the minimum weight of the working sample for purity analysis. |

For moisture analysis, the maximum time for receiving the submitted samples is 14 days after seed lot sampling.

Physiological quality: Germination test is carried out on a sample of 400 seeds in accordance with the ISTA Rules. Tests on 200 or 100 seeds are also possible depending on the need for precision. The precision of analyses is indicated in the ISTA tolerance tables.

If a germination test is requested without any specific purity analysis, pure seeds are sorted before the germination test. This analysis is not invoiced except for Grasses (*Poaceae*). This step is an integral part of the ISTA method for the evaluation of germinative faculty.

Quantity to provide for substrate checks (the retest is included in the quantities):

| | Top of paper | Rolled | Pleated paper | Sand | Organic growing media |
|----------|--------------|-----------|---------------|-------|--------------------------|
| GE-SUB-1 | 20 sheets | 12 sheets | 12 sheets | 10 kg | 8 kg |
| GE-SUB-2 | 20 sheets | 10 sheets | 10 sheets | 1 kg | 1 kg |
| GE-SUB-3 | 16 sheets | 10 sheets | 2 sheets | 1 kg | 1 kg |
| GE-SUB-4 | 96 sheets | 16 sheets | 16 sheets | 20 kg | 10 kg |

Supply of samples to the SNES



Please provide one sample per test requested with the corresponding quantity.

For OIC request, an ISTA method will be chosen if it exists.

Virology: Certain types of treatment may affect the analysis, seeds should therefore be sent untreated. If seeds has been treated with a virucidal product, please indicate this information on your order form.

Mycology:

This test is performed by detection on medium according to the following criteria:

- Without superficial disinfection for most species. If the presence of saprophytes is to high the result will be "undetermined", a new test with superficial disinfection will be proposed.
- With superficial disinfection for species that are known to have saprophytes that can compromise the analysis.

For treated seeds, a test without superficial disinfection is indicated in the price list and will be chosen.

As the method allows the detection of several pathogens simultaneously, the main pathogens are in bold in this price list and will always be indicated on the certificate. For pathogens not in bold they will be indicated on the certificate if their presence is high (> 5%) or if they were asked when the analyses were requested.

For any request for detection of other fungi, please contact SNES.

The nomenclature of fungi evolves; we therefore modify the names of pathogens to follow it. We will indicate any pathogen synonyms in brackets in the price list and test results.

In the nomenclature, "sp." means "unidentified species", "spp." means "all species" and the preceding name is the genus. If we cannot determine the species we will give as result the genus name followed by "sp.".

The denomination as sections has become obsolete, so the detection of *Fusarium*, apart from the identification (PA-ID-FUS), will be done by section classification. Some species-specific *Fusarium* will remain denominated with the species name (e.g. *F. oxysporum* on cucurbits).

Sections correspond to the classification of Nelson *and al.*; 1983, amended by Burgess *and al.*; 1994 and updated with molecular techniques (Leslie et Summerell; 2006, Carter *and al.*; 2000, Aoki et O'Donnel; 1999, Benyon *and al.*; 2000).

| Former name | Current sections | Main species |
|----------------------|-------------------------------------|---|
| | Roseum | F. avenaceum |
| Fusarium roseum | Discolor | F. culmorum, F. graminearum (Gibberella zeae), F. sambucinum, F. crookwellense |
| | Arthrosporiella | F. incarnatum (Fusarium semitectum) |
| | Sporotrichiella | F. poae, F. tricinctum (Gibberella tricincta), F. sporotrichioides, F. langsethiae |
| Fusarium sp. | Gibbosum | F. equiseti (Gibberella intricans), F. acuminatum (Gibberella acuminata) |
| Fusarium moniliforme | Liseola ou complexe G. fujikuroi | Gibberella fujikuroi (F. verticillioides, F. subglutinans), F. proliferatum |
| Fusarium oxysporum | F. elegans | F. oxysporum |
| Fusarium solani | Martiella - Ventricosum | F. solani |

Order an analysis



To SNES

| For GEVES or COFKAC certificate. | | |
|--|-----|-------|
| | | Price |
| By paper order form | | |
| Handling of the request per submitted sample and issuing of a definitive GEVES or COFRAC certificate, in French or English. | | 9.70 |
| By internet on DSN website | | |
| Handling of the request per submitted sample and issuing of a definitive GEVES or COFRAC certificate, in French or English. | | 7.40 |
| Specific handling | | |
| Handling of the request per submitted sample sent in several packaging or weighing more than 2 kg requiring the preparation of a working sample, and issuing of a definitive SNES or COFRAC certificate, in French or English. | | 41.20 |
| Supplementary certificates, specific presentation of results, priority | | |
| Duplicate certificate for adding manual singature and buffer, in French or English. | NEW | 3.10 |
| Summary table of results, or specific presentation of results. | | 31.50 |
| Raw results on .csv file (request must be entered online on DSN website). | | 0.00 |
| Priority processing, per sample. | | 18.90 |

| For an international certificate | |
|--|-------|
| 1 of all international certificate | Price |
| | PIICE |
| By paper order form | |
| Handling of each submitted sample and issuing of an Orange or Blue International Certicate, in French or English, with priority being given to the related analyses. (EC-01 + SCLI-URBI + BU-ABIODE/BU-ABIBDE) | 39.00 |
| Supplementary certificates and request for changes | |
| Provisional international certificate, in French or English. | 10.40 |
| Duplicate international certificate, in French or English. | 10.40 |
| Adding additional certificates or modification of information on an international certificate (after checking the conformity with ISTA rules). | 34.00 |

| To | Ri | a | 36 | V | EG |
|----|----------|---------|----|---|----|
| | - | [0] 4 | | w | |

Handling and results

| | Price |
|--|-------|
| Handling | |
| Handling of the sample for treated seeds. | 57.00 |
| Results | |
| Duplicates analysis certificate except photography. | 2.80 |
| New edition of result certificate. | 28.10 |
| Specific presentation of results - Contact BioGeves. | |

All Species

SEED QUALITY Physiological quality Size Duration Price Complementary determinations in addition to the germination test Detailed description of seedlings and seeds on 400 seeds. **GE-FG-DET** 1 250 41.60 Detailed description of seedlings and seeds on 200 seeds. **GE-FG-DET2** 500 20.80 Percentage of a particular type of seedling. GE-FG-PCPL 23.10 Provision of the result of repetitions. 13.40 **GE-FG-REP** Additional testing time required Additional duration of 7 days for a germination test on 400 seeds. GE-FG-7S4 1 250 16.20 Additional duration of 14 days for a germination test on 400 seeds. GE-FG-14S4 500 32.60 500 Additional duration of 7 days for a germination test on 200 seeds. GE-FG-7S2 8.20 500 Additional duration of 14 days for a germination test on 200 seeds. GE-FG-14S2 16.30 Verification of species Verification of species after germination test. **GE-ENR** 9.40 Verification of species on pelleted seeds, when only a purity test is resquested. **GE-VERIF NEW** 24.00 Tetrazolium viability test (excluding ornamental and fruit species) - For results within a week, reception of seeds on Tuesday at the latest. Tetrazolium test on 400 seeds. GE-TZ-1 500 174.00 Tetrazolium test on 200 seeds. 300 GE-TZ-2 116.00 GE-TZ-3 200 Tetrazolium test on 100 seeds. 81.00 Energy Germination energy (intermediate counting; germination capacity supplement). The date of GE-EG 500 19.90 counting for the energy varies according to the species. Cold-test on 400 seeds. GE-CO 1 250 69.00 Cold-test on 200 seeds. GE-CO2 500 44.30 Accelerated ageing of 200 seeds including germination capacity. **GE-VIEI-2** 500 90.00 Controlled deterioration of 200 seeds including germination capacity. GE-DET-1 500 90.00 Conductivity test on 200 seeds on ISTA species. **GE-CON-GLO** 500 57.00 The moisture content of seeds should be between 10 and 14 %, sample must be send in a sealed foil sachet with the indication of the water content, otherwise it would be determined by us before the test and invoiced (see test TE-SN-01). Additional cost for a conductivity test on a treated seed sample. **GE-CON-SUP** 5.40 Treatment of seeds Treatment of seeds to be performed by SNES. Seeds do not undergo fungicide treatment before **GE-TRAIT** 23.10 the germination test unless specifically requested (except for Beet). Substrate checks Determination of the water holding capacity of a substrate including moisture content. GE-SUB-1 92.00 See p.7 Determination of the pH of a substrate. GE-SUB-2 See p.7 59.00 Determination of the conductivity of a substrate. GE-SUB-3 See p.7 59.00 Assessment of the innocuity of a substrate (determination of the % of seedlings intoxicated by GE-SUB-4 See p.7 134.00 the substrate, on 2 sensitive species). GE-SUB-5 Contact SNES Viability determination of seeds in a soil or a substrate. Automated germination kinetics by image analysis Germination kinetics by image analysis (average rate of germination, kinetic curve). GE-CI **Contact SNES** Supply of detailed data on imbibition and early elongation of the root. GE-CI-4 Contact SNES Supply of seeds images during germination. GE-CI-5 **Contact SNES**

| Seed health - Prior operations | | | | |
|--|--------|------|----------|-------|
| | | Size | Duration | Price |
| Thousand Seed Weight (TSW), if not indicated on the request for bacteriology, mycology and | PA-MMS | / | / | 33.00 |
| virology tests. | | | | |

| Bacteriology - Uncoated seeds only | | | | |
|---|-------------------------|--------|-----------------------|---------------------------|
| | | Size | Duration | Price |
| Supplement fee for counting of colonies | | | | |
| 1 pathogen in 5 000 seeds. | PA-BA-19 NEW | 5 000 | | 25.00 |
| 1 pathogen in 30 000 seeds. | PA-BA-20 | 30 000 | | 61.00 |
| More than 1 pathogen in 5 000 seeds. | PA-BA-81 | 5 000 | /_ | 39.00 |
| More than 1 pathogen in 30 000 seeds. | PA-BA-82 | 30 000 | / | 114.00 |
| Mycology - See p.8 "Seed health" | | | | |
| | | Size | Duration | Price |
| Fusarium spp. | | , | | |
| Identification of <i>Fusarium</i> species in addition to detection test. | PA-ID-FUS | / | 19 days | 265.00 |
| Verticillium dahliae Agar method. | PA-ES-VERT | 400 | 19 days | 106.0 |
| Supplement for spore counting, washing methods | | | | |
| Counting by classes (0;1-10;11-100;>100). | PA-MY-DCLA | | | 64.00 |
| Counting by unit. | PA-MY-DEN | / | / | 105.00 |
| Nematology | | | | |
| | | Size | Duration | Price |
| Heterodera group schachtii, Heterodera group goettingiana, Heterodera | | | | |
| group avenae. | | | | |
| Detection and identification on soil samples. | PA-NE-SOL1 | 300 g | 30 days | 175.0 |
| Other tests | | | | |
| | | Size | Duration | Price |
| Resistance of fungal isolates to fungicides. | PA-AD-01 | | | act SNE |
| Study of the efficacy of seed disinfection/treatment products on medium or by bioassay. | PA-AD-02 PA-AD-IP | | | act SNES 50.0 0 |
| Identification of pathogens isolated and provided on medium - Supply 2 boxes/isolates. Isolation of strains from symptoms. | PA-AD-IP PA-ISOLEM | | 19 days | 50.00 |
| Isolation of strains from seeds. | PA-ISOSEM | | | 107.0 |
| Identification of pathogens on plant material. | PA-DI-PEC | | Cont | act SNE |
| Feasibility on a case-by-case basis. Prices below are indicated for information. They will be | | | | |
| charged depending on the observed symptoms. Handling of the sample. | PA-DI-PEC | | | 57.0 |
| Identification based on symptoms. | PA-DI-PEC PA-DI-MICR | | | 98.0 |
| Mycological identification after incubation. | PA-DI-MY | | | 200.0 |
| Bacteriological identification after incubation. | PA-DI-BA | | | 100.0 |
| Confirmation by pathogenicity test. | PA-DI-PP | | | 122.0 |
| Virological identification by immunological test. | PA-DI-ELIS | | | 215.0 |
| Virological identification virologic by biotest. | PA-DI-IND | | | 69.0 |
| Analytical Profile Index (API). | PA-DI-API | | / | 190.0 |
| PCR. | PA-DI-PCR | / | / | 121.0 |
| EVALUATION OF VARIETIES | | | | |
| Determination of the identity and the varietal purity | | | | |
| · | | Size | Duration | Price |
| Standard protocol. | SEV-CV | | | 345.00 |
| Specific study. | SEV-CV1 | | Cor | ntact SEV |
| Genotyping by molecular biology | | | | |
| Variabel identify, annual, CCD | DI C DE4 CCD C'C 4 | Size | Duration Contact F | Price |
| Varietal identity control - SSR. | BI-G-BM-SSR-CID-1 | | Contact E | DIOGENES |
| Varietal comparison - SSR. | BI-G-BM-SSR-COMP | | Contact E |):aCE\ /EC |

All Species

| | | Size | Duration | Price |
|---|--------------------|------|-----------|---------|
| Genetic purity analysis - SSR - 180 seeds. | BI-G-BM-SSR-PU-180 | | Contact B | ioGEVES |
| Genetic purity analysis - SSR - 8 x 10 seeds. | BI-G-BM-SSR-PUR-10 | | Contact B | ioGEVES |
| Seed mixture detection. | BI-G-BM-SSR-PUR-40 | | Contact B | ioGEVES |
| Varietal purity analysis - SSR - 90 seeds. | BI-G-BM-SSR-PUR-90 | | Contact B | ioGEVES |
| Varietal description - SSR. | BI-G-BM-SSR-DVAR | | Contact B | ioGEVES |
| DNA extraction. | BI-G-BM-EXT | | Contact B | ioGEVES |
| Varietal identity control - SNP. | BI-G-BM-SNP-CID | | Contact B | ioGEVES |
| Hybrid Conformity - SNP. | BI-G-BM-SNP-CONF | | Contact B | ioGEVES |
| Varietal comparison - SNP. | BI-G-BM-SNP-COMP | | Contact B | ioGEVES |
| Genetic purity analysis - SNP. | BI-G-BM-SNP-PUR | | Contact B | ioGEVES |
| Varietal description - SNP. | BI-G-BM-SNP-DVAR | | Contact B | ioGEVES |
| Standardization of DNA concentration & distribution in plate. | BI-G-CUST-GEN-3 | | Contact B | ioGEVES |
| Analysis of genetic diversity. | BI-G-CUST-GEN-2 | | Contact B | ioGEVES |
| Migration run - Capillary sequencer - plate. | BI-G-BM-RUN | | Contact B | ioGEVES |
| DNA assay. | BI-G-BM-DOS | | Contact B | ioGEVES |
| Development of genotyping method. | BI-G-METH | | Contact B | ioGEVES |
| Customised genotyping. | BI-G-CUST | | Contact B | ioGEVES |

| Technological quality: biochemical tests | | | | |
|--|--------------------|------|------------|--------|
| | | Size | Duration | Price |
| SPEC - custom analysis. | BI-B-CUST-DEV-SPEC | | Contact Bi | oGEVES |
| RMN - custom analysis. | BI-B-CUST-DEV-RMN | | Contact Bi | oGEVES |
| CPG - custom analysis. | BI-B-CUST-DEV-CPG | | Contact Bi | oGEVES |
| NIRS - custom analysis. | BI-B-CUST-DEV-NIRS | | Contact Bi | oGEVES |
| HPLC - custom analysis. | BI-B-CUST-DEV-HPLC | | Contact Bi | oGEVES |
| Tannin content (assay by spectrophotometry). | BI-B-SPEC-TAN-GEN | | Contact Bi | oGEVES |
| Fatty acid composition. | BI-B-CPG-AG-GEN | | Contact Bi | oGEVES |
| Glucosinolate content (HPLC). | BI-B-HPLC-GLU-GEN | | Contact Bi | oGEVES |
| Antitrypsic activity. | BI-B-SPECT-FAT-GEN | | Contact Bi | oGEVES |
| Glucosinolate content (NIRS). | BI-B-NIRS-NGLS | | Contact Bi | oGEVES |
| Spectrochlorophyll. | BI-B-SPEC-CHLO | | Contact Bi | oGEVES |
| Customised biochemical molecule assays (NIRS model development, analytical chemistry). | BI-B-CUST | | Contact Bi | oGEVES |
| Oil content (NMR). | BI-B-RMN-H | | Contact Bi | oGEVES |
| Water content (NMR). | BI-B-RMN-E | | Contact Bi | oGEVES |
| Phytates by spectrophotometry. | BI-B-SPEC-PHY | NEW | Contact Bi | oGEVES |

| Other tests | | | | |
|----------------------------------|--------------|------|------------|--------|
| | | Size | Duration | Price |
| WDV virus detection test by PCR. | BI-D-VIR-WDV | | Contact Bi | oGEVES |

Annual subscription to the variety denomination class test

Price

| All species - 10 tests. | SEV-DENOS-10 | 215.00 |
|--------------------------|---------------|---------|
| All species - 20 tests. | SEV-DENOS-20 | 405.00 |
| All species - 50 tests. | SEV-DENOS-50 | 950.00 |
| All species - 100 tests. | SEV-DENOS-100 | 1830.00 |
| All species - 200 tests. | SEV-DENOS-200 | 3580.00 |

PUBLICATIONS (Contact SNES)

Technical sheet for analysis of specific purity and counting of all other seeds

Purity and determination of other seeds by number: methodology.

AP-M-1

All Species

| Identification data sheet of seeds and other impurities | |
|---|-----------|
| Echinochloa crus-galli, Echinochloa colona, Panicum capillare, Panicum maximum, Setaria pumila, Setaria veridis. | AP-A-01 |
| Avena fatua-Avena sativa. | AP-A-02 |
| Germination analysis method sheet | |
| Germination method of different species. | GE-M-ESP |
| Identification data sheet of seeds and other impurities | |
| Polygonaceae (Persicaria maculosa, Persicaria lapathifolia, Fallopia convolvulus, Polygonum aviculare, Rumex sp., Rumex acetosella, Rumex maritimus). | AP-A-03 |
| Chenopodium sp., Atriplex sp., Amaranthus sp., Reseda sp., Myosotis sp. | AP-A-04 |
| Asteraceae (Anthemis arvensis, Glebionis segetum, Chicorium sp., Tripleurospermum inodorum, Helminthotheca echioïdes, Lapsana communis, Lactuca sativa, Sonchus spp., Cirsium arvense, Cirsium vulgare, Centaurea cyanus). | AP-A-06 |
| Cuscuta spp. | AP-P-1 |
| Claviceps purpurea - Sclerotinia sclerotiorum. | AP-P-2 |
| Self-control kit | |
| On request, components are sent separately accompanied with an instructional material. Contact SNES. | KIT-AUTO |
| I.D.Seed® On-line picture library, an aid to the identification of seeds - In French | |
| I.D.Seed® - Complete collection. Resgistration on http://mediatheque.geves.fr | IDSEED-1 |
| Identification data sheet of fungal pathogens | |
| Altenaria linariae, A. alternata, A. brassicae, A. brassicicola, A. cucumerina, A. dauci, A. japonica, A. linicola, A. padwickii, A. petroselini, Alternariaster helianthi, Ascochyta medicaginicola, Bipolaris oryzae, Botryotinia squamosa, Botrytis cinerea, Ciborinia allii, Colletotrichum graminicola, C. truncatum, Complexe Phomopsis, Didymella pisi, Exserohilum turcicum, Itersonilia perplexans, Phomopsis helianthi, Sarocladium strictum, Sclerotinia sclerotiorum. | PA-T-PATH |
| Identification data sheet of nematodes | |
| Ditylenchus dipsaci, D. destructor, Aphelenchoides besseyi, A. fragariae. | PA-T-NEM |
| Identification data sheet of fungal saprophytes | |
| Sheet containing the main fungal saprophytes present in analysis on media. | PA-T-SAPR |

Seed mixture species

SEED QUALITY

Physical quality

| Size | Duration | Price |
|------|----------|-------|
| | | |

Purity analysis test and determination of the composition of a seed mixture of species Only on naked seeds

| Less than 4 components WITH declared composition ² . | PU-MEL-01 | / | 60 days | 513.00 |
|--|------------|---|---------|----------|
| From 4 components WITH declared composition ² . | PU-MEL-02 | | Cont | act SNES |
| WITHOUT declared composition. | PU-MEL-03 | / | 60 days | 843.00 |
| Preparation of pure seed for germination testing | | | | |
| Seed mixture (less than 4 components) WITH declared composition ² . | PU-PR-19 | / | / | 212.00 |
| From 4 components WITH declared composition ² . | PU-PR-22 | | Cont | act SNES |
| WITHOUT declared composition. | PU-PR-19-1 | / | / | 508.00 |
| Preparation of pure seeds in dragees on coated seed mixture. | PU-PR-19-2 | | / | 35.90 |

 $^{^{\}rm 2}$ Provide the % of species in the seed mixture.

Physiological quality ³

Germination test on 400 seeds


Species mixture by component. All the species of the seed mixture will be analyzed whatever is the proportion, except opposite request.

Germination test on 200 seeds

Species mixture by component. All the species of the seed mixture will be analyzed whatever is the proportion, except opposite request.

GE-FG-19-2


³ See details of price and size in the chapter of the species.




| Emilia do unitariligi di uni danci decud | | | | |
|--|--------------------------|----------------|--------------------|----------------------------------|
| Searching of 1 to $\overline{4}$ species (except for <i>Orobanchaceae</i>). Indicate the name of the species to be searched. | SP-LI-01 | ISTA weight | / | 64.00 |
| Searching of 5 to 8 species (except for <i>Orobanchaceae</i>). Indicate the name of the species to be searched. | SP-LI-02 | ISTA weight | / | 102.00 |
| Searching of more than 8 species (except for <i>Orobanchaceae</i>). Indicate the name of the species to be searched. | SP-LI-20 | | Cont | act SNES |
| Searching of <i>Orobanche</i> sp. Only on UNTREATED and UNCOATED seeds. Analyse performed on a separate, sealed, submitted subsample. | SP-ORO | ISTA weight | / | 75.00 |
| Searching of <i>Striga</i> sp. Only on UNTREATED and UNCOATED seeds. Analyse performed on a separate, sealed, submitted subsample. | SP-STRIGA | ISTA weight | / | 75.00 |
| Searching of <i>Orobanche</i> sp. and <i>Striga</i> sp. Only on UNTREATED and UNCOATED seeds. Analyse performed on a separate, sealed, submitted subsample. | SP-ORO-STR | ISTA weight | / | 110.00 |
| Tests on coated seeds | | | | |
| Purity on coated seeds. | PU-IS-21 | 2 500 | / | 34.30 |
| Pelleting material removal and full counting on 2 500 coated seeds. Only on UNTREATED seeds. | SP-ENR2500 | 2 500 | / | 102.00 |
| Pelleting material removal and full counting on 7 500 coated seeds. Only on UNTREATED seeds. | SP-ENR-TOT | 7 500 | / | 309.00 |
| Pelleting material removal and limited counting of other seeds from 1 to 3 botanical species, on 7 500 coated seeds. Only on UNTREATED seeds. | SP-ENR-LIM | 7 500 | / | 242.00 |
| Moisture content - Provide seeds in watertight bags from which as much air as possible has been extracted | | | | |
| | | | | 20.70 |
| Oven method. | TE-SN-01 | ISTA weight | / | 20.70 |
| | TE-SN-01 | ISTA weight | / | 20.70 |
| Oven method. Identification of individual seeds Visual identification by species. | TE-SN-01 | ISTA weight | / | 20.70 34.60 |
| Identification of individual seeds | | ISTA weight | / | |
| Identification of individual seeds Visual identification by species. | | ISTA weight / | / Duration | 34.60 |
| Identification of individual seeds Visual identification by species. | | / | / / Duration | 34.60 |
| Identification of individual seeds Visual identification by species. Physiological quality | | / | / Duration | 34.60 Price |
| Identification of individual seeds Visual identification by species. Physiological quality Germination test on 400 seeds | ID-IS-01 | / Size | Duration / | 34.60 Price 65.00 |
| Identification of individual seeds Visual identification by species. Physiological quality Germination test on 400 seeds Vegetables (excluding corn salad). Corn salad. The germination capacity tests of corn salad seeds are carried out using several methods on 400 seeds: 2 methods with and without sodium hypochlorite disinfection from January 1st to May 31st and 2 methods with and without sodium hypochlorite disinfection and gibberellin from June | ID-IS-01 GE-FG-18-4 | Size | Duration / | |
| Identification of individual seeds Visual identification by species. Physiological quality Germination test on 400 seeds Vegetables (excluding corn salad). Corn salad. The germination capacity tests of corn salad seeds are carried out using several methods on 400 seeds: 2 methods with and without sodium hypochlorite disinfection from January 1st to May 31st and 2 methods with and without sodium hypochlorite disinfection and gibberellin from June 1st to December 31st. | ID-IS-01 GE-FG-18-4 | Size | Duration / | 34.60 Price 65.00 66.00 |
| Identification of individual seeds Visual identification by species. Physiological quality Germination test on 400 seeds Vegetables (excluding corn salad). Corn salad. The germination capacity tests of corn salad seeds are carried out using several methods on 400 seeds: 2 methods with and without sodium hypochlorite disinfection from January 1st to May 31st and 2 methods with and without sodium hypochlorite disinfection and gibberellin from June 1st to December 31st. Germination test on 200 seeds | GE-FG-18-4 GE-FG-22-4 | 1 250 1 250 | Duration / | 34.60 Price 65.00 |


| | | Size | Duration | Pric |
|--|--|---|---|--|
| | | | | |
| Additional | CE EC HAD | , | , | 7.0 |
| Additional cost for manual sowing of fragile seeds of bean. | GE-FG-HAR | | | |
| Germination tests on bulbs and bulblets On 400 seeds. | GE-BULB-4 | 1 250 | , | 151. |
| On 200 seeds. | GE-BULB-2 | 500 | | 122. |
| | GL-DOLD-2 | 300 | | 122. |
| .ettuce specific cold-test On 400 seeds. | GE-EGFG-4 | 1 250 | , | 92. |
| On 200 seeds. | GE-EGFG-2 | 500 | | 54 |
| /erification of species | | | | |
| /erification of species after germination test. | GE-ENR | / | / | 9 |
| | OL LIVI | , , , , , , , , , , , , , , , , , , , | | |
| /igour tests Conductivity test on 200 seeds on ISTA species. | GE-CON-GLO | 500 | / | 57 |
| The moisture content of seeds should be between 10 and 14 %, sample must be send in a sealed | GE-CON-GEO | 300 | , | 3, |
| oil sachet with the indication of the water content, otherwise it would be determined by us | | | | |
| pefore the test and invoiced (see test TE-SN-01). | | | | |
| Determination of the rate of usable Tomato plants | | | | |
| On 400 seeds. | GE-TX-PL-2 | 500 | | 105 |
| On 200 seeds. | GE-TX-PL-1 | 300 | / | 80 |
| Treatment of seeds | | | | |
| reatment of seeds to be performed by SNES. Seeds do not undergo fungicide treatment before | GE-TRAIT | / | / | 23 |
| he germination test unless specifically requested (except for Beet). | | | | |
| Destruistant Hussahad anada subs | | | | |
| Bacteriology - Uncoated seeds only | | | | |
| | | | | |
| | | Size | Duration | Pı |
| Eggplant, Pepper, Tomato - Detection of 1 pathogen | | Size | Duration | Pı |
| Eggplant, Pepper, Tomato - Detection of 1 pathogen Pseudomonas syringae pv. tomato (Pst) | | Size | Duration | Pı |
| | PA-BA-25 | Size 30 000 | Duration 36 days | Pr 219 |
| Pseudomonas syringae pv. tomato (Pst) | PA-BA-25 | | | |
| Pseudomonas syringae pv. tomato (Pst) Agar method + pathogenicity test in case of suspect colonies. | PA-BA-25 PA-BA-26 | | | 219 |
| Pseudomonas syringae pv. tomato (Pst) Agar method + pathogenicity test in case of suspect colonies. Kanthomonas spp. pathogenic on Tomato and Pepper (X spp) Agar method + identification of strains by PCR in case of suspect colonies (ISF). Agar method + identification of strains by PCR in case of suspect colonies (ISF). | | 30 000 | 36 days | 219 245 |
| Pseudomonas syringae pv. tomato (Pst) Agar method + pathogenicity test in case of suspect colonies. Kanthomonas spp. pathogenic on Tomato and Pepper (X spp) Agar method + identification of strains by PCR in case of suspect colonies (ISF). | PA-BA-26 | 30 000 30 000 | 36 days | 219 |
| Pseudomonas syringae pv. tomato (Pst) Agar method + pathogenicity test in case of suspect colonies. Kanthomonas spp. pathogenic on Tomato and Pepper (X spp) Agar method + identification of strains by PCR in case of suspect colonies (ISF). Agar method + identification of strains by PCR in case of suspect colonies (ISF). | PA-BA-26 | 30 000 30 000 | 36 days | 219 245 |
| Pseudomonas syringae pv. tomato (Pst) Agar method + pathogenicity test in case of suspect colonies. Kanthomonas spp. pathogenic on Tomato and Pepper (X spp) Agar method + identification of strains by PCR in case of suspect colonies (ISF). Agar method + identification of strains by PCR in case of suspect colonies (ISF). Analyse done with 1 subsample on 10 000 seeds maximum. Pseudomonas corrugata (Pc) Agar method + pathogenicity test in case of suspect colonies. | PA-BA-26 | 30 000 30 000 | 36 days | 219 245 170 |
| Pseudomonas syringae pv. tomato (Pst) Agar method + pathogenicity test in case of suspect colonies. Kanthomonas spp. pathogenic on Tomato and Pepper (X spp) Agar method + identification of strains by PCR in case of suspect colonies (ISF). Agar method + identification of strains by PCR in case of suspect colonies (ISF). Analyse done with 1 subsample on 10 000 seeds maximum. | PA-BA-26 PA-BA-26-1 | 30 000 30 000 Max 10 000 | 36 days 34 days 34 days | 219 245 170 |
| Pseudomonas syringae pv. tomato (Pst) Agar method + pathogenicity test in case of suspect colonies. Kanthomonas spp. pathogenic on Tomato and Pepper (X spp) Agar method + identification of strains by PCR in case of suspect colonies (ISF). Agar method + identification of strains by PCR in case of suspect colonies (ISF). Analyse done with 1 subsample on 10 000 seeds maximum. Pseudomonas corrugata (Pc) Agar method + pathogenicity test in case of suspect colonies. | PA-BA-26 PA-BA-26-1 | 30 000 30 000 Max 10 000 | 36 days 34 days 34 days | 219 245 170 |
| Pseudomonas syringae pv. tomato (Pst) Agar method + pathogenicity test in case of suspect colonies. Kanthomonas spp. pathogenic on Tomato and Pepper (X spp) Agar method + identification of strains by PCR in case of suspect colonies (ISF). Analyse done with 1 subsample on 10 000 seeds maximum. Pseudomonas corrugata (Pc) Agar method + pathogenicity test in case of suspect colonies. Eggplant, Pepper, Tomato - Detection of 2 pathogens Pst + X spp Agar method + pathogenicity test and/or identification of strains by PCR in case of suspect | PA-BA-26 PA-BA-26-1 | 30 000 30 000 Max 10 000 | 36 days 34 days 34 days | 219 245 170 295 |
| Pseudomonas syringae pv. tomato (Pst) Agar method + pathogenicity test in case of suspect colonies. Kanthomonas spp. pathogenic on Tomato and Pepper (X spp) Agar method + identification of strains by PCR in case of suspect colonies (ISF). Agar method + identification of strains by PCR in case of suspect colonies (ISF). Analyse done with 1 subsample on 10 000 seeds maximum. Pseudomonas corrugata (Pc) Agar method + pathogenicity test in case of suspect colonies. Eggplant, Pepper, Tomato - Detection of 2 pathogens Pst + X spp | PA-BA-26-1 PA-BA-26-1 PA-BA-92 | 30 000 30 000 Max 10 000 30 000 | 36 days 34 days 34 days 36 days | 219 245 170 295 |
| Pseudomonas syringae pv. tomato (Pst) Agar method + pathogenicity test in case of suspect colonies. Kanthomonas spp. pathogenic on Tomato and Pepper (X spp) Agar method + identification of strains by PCR in case of suspect colonies (ISF). Agar method + identification of strains by PCR in case of suspect colonies (ISF). Analyse done with 1 subsample on 10 000 seeds maximum. Pseudomonas corrugata (Pc) Agar method + pathogenicity test in case of suspect colonies. Eggplant, Pepper, Tomato - Detection of 2 pathogens Pst + X spp Agar method + pathogenicity test and/or identification of strains by PCR in case of suspect colonies (ISF for Xanthomonas). Pst + Pc | PA-BA-26 PA-BA-26-1 PA-BA-92 PA-BA-40 | 30 000 30 000 Max 10 000 30 000 | 36 days 34 days 34 days 36 days | 219 245 170 295 |
| Agar method + pathogenicity test in case of suspect colonies. Kanthomonas spp. pathogenic on Tomato and Pepper (X spp) Agar method + identification of strains by PCR in case of suspect colonies (ISF). Agar method + identification of strains by PCR in case of suspect colonies (ISF). Analyse done with 1 subsample on 10 000 seeds maximum. Pseudomonas corrugata (Pc) Agar method + pathogenicity test in case of suspect colonies. Eggplant, Pepper, Tomato - Detection of 2 pathogens Pst + X spp Agar method + pathogenicity test and/or identification of strains by PCR in case of suspect colonies (ISF for Xanthomonas). Pst + Pc Agar method + identification of strains by pathogenicity test in case of suspect colonies. | PA-BA-26-1 PA-BA-26-1 PA-BA-92 | 30 000 30 000 Max 10 000 30 000 | 36 days 34 days 34 days 36 days | 219 245 170 295 |
| Pseudomonas syringae pv. tomato (Pst) Agar method + pathogenicity test in case of suspect colonies. Kanthomonas spp. pathogenic on Tomato and Pepper (X spp) Agar method + identification of strains by PCR in case of suspect colonies (ISF). Agar method + identification of strains by PCR in case of suspect colonies (ISF). Analyse done with 1 subsample on 10 000 seeds maximum. Pseudomonas corrugata (Pc) Agar method + pathogenicity test in case of suspect colonies. Eggplant, Pepper, Tomato - Detection of 2 pathogens Pst + X spp Agar method + pathogenicity test and/or identification of strains by PCR in case of suspect colonies (ISF for Xanthomonas). Pst + Pc | PA-BA-26 PA-BA-26-1 PA-BA-92 PA-BA-40 | 30 000 30 000 Max 10 000 30 000 | 36 days 34 days 36 days 40 days | 219 245 170 295 |
| Pseudomonas syringae pv. tomato (Pst) Agar method + pathogenicity test in case of suspect colonies. Kanthomonas spp. pathogenic on Tomato and Pepper (X spp) Agar method + identification of strains by PCR in case of suspect colonies (ISF). Agar method + identification of strains by PCR in case of suspect colonies (ISF). Analyse done with 1 subsample on 10 000 seeds maximum. Pseudomonas corrugata (Pc) Agar method + pathogenicity test in case of suspect colonies. Eggplant, Pepper, Tomato - Detection of 2 pathogens Pst + X spp Agar method + pathogenicity test and/or identification of strains by PCR in case of suspect colonies (ISF for Xanthomonas). Pst + Pc Agar method + identification of strains by pathogenicity test in case of suspect colonies. Eggplant, Pepper, Tomato - Supplement fee pathogenicity test | PA-BA-26 PA-BA-26-1 PA-BA-92 PA-BA-40 | 30 000 30 000 Max 10 000 30 000 | 36 days 34 days 36 days 40 days | 219 245 170 295 |
| Pseudomonas syringae pv. tomato (Pst) Agar method + pathogenicity test in case of suspect colonies. Kanthomonas spp. pathogenic on Tomato and Pepper (X spp) Agar method + identification of strains by PCR in case of suspect colonies (ISF). Analyse done with 1 subsample on 10 000 seeds maximum. Pseudomonas corrugata (Pc) Agar method + pathogenicity test in case of suspect colonies. Eggplant, Pepper, Tomato - Detection of 2 pathogens Pst + X spp Agar method + pathogenicity test and/or identification of strains by PCR in case of suspect colonies (ISF for Xanthomonas). Pst + Pc Agar method + identification of strains by pathogenicity test in case of suspect colonies. Eggplant, Pepper, Tomato - Supplement fee pathogenicity test K spp Confirmation by pathogenicity test of PCR positive isolates. | PA-BA-26 PA-BA-26-1 PA-BA-92 PA-BA-40 | 30 000 30 000 Max 10 000 30 000 | 36 days 34 days 36 days 40 days | 219 245 170 295 346 |
| Pseudomonas syringae pv. tomato (Pst) Agar method + pathogenicity test in case of suspect colonies. Kanthomonas spp. pathogenic on Tomato and Pepper (X spp) Agar method + identification of strains by PCR in case of suspect colonies (ISF). Analyse done with 1 subsample on 10 000 seeds maximum. Pseudomonas corrugata (Pc) Agar method + pathogenicity test in case of suspect colonies. Eggplant, Pepper, Tomato - Detection of 2 pathogens Pst + X spp Agar method + pathogenicity test and/or identification of strains by PCR in case of suspect colonies (ISF for Xanthomonas). Pst + Pc Agar method + identification of strains by pathogenicity test in case of suspect colonies. Eggplant, Pepper, Tomato - Supplement fee pathogenicity test (S spp Confirmation by pathogenicity test of PCR positive isolates. Eggplant, Pepper - Detection of 1 pathogen | PA-BA-26-1 PA-BA-92 PA-BA-40 PA-BA-127 | 30 000 30 000 Max 10 000 30 000 | 36 days 34 days 34 days 40 days | |
| Pseudomonas syringae pv. tomato (Pst) Agar method + pathogenicity test in case of suspect colonies. Kanthomonas spp. pathogenic on Tomato and Pepper (X spp) Agar method + identification of strains by PCR in case of suspect colonies (ISF). Agar method + identification of strains by PCR in case of suspect colonies (ISF). Analyse done with 1 subsample on 10 000 seeds maximum. Pseudomonas corrugata (Pc) Agar method + pathogenicity test in case of suspect colonies. Eggplant, Pepper, Tomato - Detection of 2 pathogens Pst + X spp Agar method + pathogenicity test and/or identification of strains by PCR in case of suspect colonies (ISF for Xanthomonas). Pst + Pc Agar method + identification of strains by pathogenicity test in case of suspect colonies. Eggplant, Pepper, Tomato - Supplement fee pathogenicity test K spp Confirmation by pathogenicity test of PCR positive isolates. Eggplant, Pepper - Detection of 1 pathogen Clavibacter michiganensis subsp. michiganensis (Cmm) | PA-BA-26 PA-BA-26-1 PA-BA-92 PA-BA-40 PA-BA-127 PA-PP-XPP | 30 000 30 000 Max 10 000 30 000 30 000 | 36 days 34 days 36 days 40 days 10 days | 219 245 170 295 346 299 |
| Reseudomonas syringae pv. tomato (Pst) Agar method + pathogenicity test in case of suspect colonies. Reanthomonas spp. pathogenic on Tomato and Pepper (X spp) Agar method + identification of strains by PCR in case of suspect colonies (ISF). Agar method + identification of strains by PCR in case of suspect colonies (ISF). Agar method + identification of strains by PCR in case of suspect colonies (ISF). Agar method + pathogenicity test in case of suspect colonies. Agar method + pathogenicity test in case of suspect colonies. Agar method + pathogenicity test and/or identification of strains by PCR in case of suspect colonies (ISF for Xanthomonas). Agar method + pathogenicity test and/or identification of strains by PCR in case of suspect colonies (ISF for Xanthomonas). Agar method + identification of strains by pathogenicity test in case of suspect colonies. Agar method + identification of strains by pathogenicity test in case of suspect colonies. Agar method + pathogenicity test of PCR positive isolates. Agaplant, Pepper - Detection of 1 pathogen Clavibacter michiganensis subsp. michiganensis (Cmm) Agar method. (M-GEVES/SV/MO/006). | PA-BA-26 PA-BA-26-1 PA-BA-92 PA-BA-40 PA-BA-127 PA-PP-XPP | 30 000 30 000 Max 10 000 30 000 30 000 / NEW 30 000 | 36 days 34 days 36 days 40 days 10 days | 219 245 170 295 346 299 |
| Pseudomonas syringae pv. tomato (Pst) Agar method + pathogenicity test in case of suspect colonies. (anthomonas spp. pathogenic on Tomato and Pepper (X spp) Agar method + identification of strains by PCR in case of suspect colonies (ISF). Agar method + identification of strains by PCR in case of suspect colonies (ISF). Analyse done with 1 subsample on 10 000 seeds maximum. Pseudomonas corrugata (Pc) Agar method + pathogenicity test in case of suspect colonies. Eggplant, Pepper, Tomato - Detection of 2 pathogens Pst + X spp Agar method + pathogenicity test and/or identification of strains by PCR in case of suspect colonies (ISF for Xanthomonas). Pst + Pc Agar method + identification of strains by pathogenicity test in case of suspect colonies. Eggplant, Pepper, Tomato - Supplement fee pathogenicity test (Spp Confirmation by pathogenicity test of PCR positive isolates. Eggplant, Pepper - Detection of 1 pathogen Clavibacter michiganensis subsp. michiganensis (Cmm) Agar method. (M-GEVES/SV/MO/006). Agar method. (M-GEVES/SV/MO/006). Analyse done with 1 subsample on 10 000 seeds | PA-BA-26 PA-BA-26-1 PA-BA-92 PA-BA-40 PA-BA-127 PA-PP-XPP | 30 000 30 000 Max 10 000 30 000 30 000 | 36 days 34 days 36 days 40 days 10 days | 219 245 170 295 346 299 |
| Pseudomonas syringae pv. tomato (Pst) Agar method + pathogenicity test in case of suspect colonies. Kanthomonas spp. pathogenic on Tomato and Pepper (X spp) Agar method + identification of strains by PCR in case of suspect colonies (ISF). Analyse done with 1 subsample on 10 000 seeds maximum. Pseudomonas corrugata (Pc) Agar method + pathogenicity test in case of suspect colonies. Eggplant, Pepper, Tomato - Detection of 2 pathogens Pst + X spp Agar method + pathogenicity test and/or identification of strains by PCR in case of suspect colonies (ISF for Xanthomonas). Pst + Pc Agar method + identification of strains by pathogenicity test in case of suspect colonies (ISF for Xanthomonas). Pst + Pc Agar method + identification of strains by pathogenicity test in case of suspect colonies. Eggplant, Pepper, Tomato - Supplement fee pathogenicity test (S spp Confirmation by pathogenicity test of PCR positive isolates. Eggplant, Pepper - Detection of 1 pathogen Clavibacter michiganensis subsp. michiganensis (Cmm) Agar method. (M-GEVES/SV/MO/006). Agar method. (M-GEVES/SV/MO/006). Analyse done with 1 subsample on 10 000 seeds maximum. | PA-BA-26 PA-BA-26-1 PA-BA-92 PA-BA-40 PA-BA-127 PA-PP-XPP | 30 000 30 000 Max 10 000 30 000 30 000 / NEW 30 000 | 36 days 34 days 36 days 40 days 10 days | 219 245 170 295 346 299 |
| Pseudomonas syringae pv. tomato (Pst) Agar method + pathogenicity test in case of suspect colonies. Kanthomonas spp. pathogenic on Tomato and Pepper (X spp) Agar method + identification of strains by PCR in case of suspect colonies (ISF). Agar method + identification of strains by PCR in case of suspect colonies (ISF). Analyse done with 1 subsample on 10 000 seeds maximum. Pseudomonas corrugata (Pc) Agar method + pathogenicity test in case of suspect colonies. Eggplant, Pepper, Tomato - Detection of 2 pathogens Pst + X spp Agar method + pathogenicity test and/or identification of strains by PCR in case of suspect colonies (ISF for Xanthomonas). Pst + Pc Agar method + identification of strains by pathogenicity test in case of suspect colonies. Eggplant, Pepper, Tomato - Supplement fee pathogenicity test K spp Confirmation by pathogenicity test of PCR positive isolates. Eggplant, Pepper - Detection of 1 pathogen Clavibacter michiganensis subsp. michiganensis (Cmm) Agar method. (M-GEVES/SV/MO/006). Agar method. (M-GEVES/SV/MO/006). Analyse done with 1 subsample on 10 000 seeds maximum. Eggplant, Pepper - Detection of 2 pathogens | PA-BA-26 PA-BA-26-1 PA-BA-92 PA-BA-40 PA-BA-127 PA-PP-XPP | 30 000 30 000 Max 10 000 30 000 30 000 / NEW 30 000 | 36 days 34 days 36 days 40 days 10 days | 219 245 170 295 346 299 |
| Agar method + pathogenicity test in case of suspect colonies. Kanthomonas spp. pathogenic on Tomato and Pepper (X spp) Agar method + identification of strains by PCR in case of suspect colonies (ISF). Agar method + identification of strains by PCR in case of suspect colonies (ISF). Agar method + identification of strains by PCR in case of suspect colonies (ISF). Analyse done with 1 subsample on 10 000 seeds maximum. Pseudomonas corrugata (Pc) Agar method + pathogenicity test in case of suspect colonies. Eggplant, Pepper, Tomato - Detection of 2 pathogens PSt + X spp Agar method + pathogenicity test and/or identification of strains by PCR in case of suspect colonies (ISF for Xanthomonas). PSt + Pc Agar method + identification of strains by pathogenicity test in case of suspect colonies. Eggplant, Pepper, Tomato - Supplement fee pathogenicity test K spp Confirmation by pathogenicity test of PCR positive isolates. Eggplant, Pepper - Detection of 1 pathogen Clavibacter michiganensis subsp. michiganensis (Cmm) Agar method. (M-GEVES/SV/MO/006). Agar method. (M-GEVES/SV/MO/006). Analyse done with 1 subsample on 10 000 seeds maximum. Eggplant, Pepper - Detection of 2 pathogens Cmm + X spp | PA-BA-26-1 PA-BA-92 PA-BA-40 PA-BA-127 PA-PP-XPP PA-BA-23-A PA-BA-23-B | 30 000 30 000 Max 10 000 30 000 30 000 / NEW 30 000 NEW Max 10 000 | 36 days 34 days 34 days 40 days 40 days 10 days 33 days 33 days | 219 245 170 295 346 299 76 352 228 |
| Pseudomonas syringae pv. tomato (Pst) Agar method + pathogenicity test in case of suspect colonies. Kanthomonas spp. pathogenic on Tomato and Pepper (X spp) Agar method + identification of strains by PCR in case of suspect colonies (ISF). Analyse done with 1 subsample on 10 000 seeds maximum. Pseudomonas corrugata (Pc) Agar method + pathogenicity test in case of suspect colonies. Eggplant, Pepper, Tomato - Detection of 2 pathogens Pst + X spp Agar method + pathogenicity test and/or identification of strains by PCR in case of suspect colonies (ISF for Xanthomonas). Pst + Pc Agar method + identification of strains by pathogenicity test in case of suspect colonies (ISF for Xanthomonas). Pst + Pc Agar method + identification of strains by pathogenicity test in case of suspect colonies. Eggplant, Pepper, Tomato - Supplement fee pathogenicity test (S spp Confirmation by pathogenicity test of PCR positive isolates. Eggplant, Pepper - Detection of 1 pathogen Clavibacter michiganensis subsp. michiganensis (Cmm) Agar method. (M-GEVES/SV/MO/006). Agar method. (M-GEVES/SV/MO/006). Analyse done with 1 subsample on 10 000 seeds maximum. | PA-BA-26 PA-BA-26-1 PA-BA-92 PA-BA-40 PA-BA-127 PA-PP-XPP | 30 000 30 000 Max 10 000 30 000 30 000 / NEW 30 000 NEW Max 10 000 | 36 days 34 days 36 days 40 days 10 days | 219 245 170 295 346 299 |
| Pseudomonas syringae pv. tomato (Pst) Agar method + pathogenicity test in case of suspect colonies. Kanthomonas spp. pathogenic on Tomato and Pepper (X spp) Agar method + identification of strains by PCR in case of suspect colonies (ISF). Analyse done with 1 subsample on 10 000 seeds maximum. Pseudomonas corrugata (Pc) Agar method + pathogenicity test in case of suspect colonies. Eggplant, Pepper, Tomato - Detection of 2 pathogens Pst + X spp Agar method + pathogenicity test and/or identification of strains by PCR in case of suspect colonies (ISF for Xanthomonas). Pst + Pc Agar method + identification of strains by pathogenicity test in case of suspect colonies. Eggplant, Pepper, Tomato - Supplement fee pathogenicity test K spp Confirmation by pathogenicity test of PCR positive isolates. Eggplant, Pepper - Detection of 1 pathogen Clavibacter michiganensis subsp. michiganensis (Cmm) Agar method. (M-GEVES/SV/MO/006). Agar method. (M-GEVES/SV/MO/006). Analyse done with 1 subsample on 10 000 seeds maximum. Eggplant, Pepper - Detection of 2 pathogens Cmm + X spp Agar method + identification of strains by PCR and/or pathogenicity test in case of suspect | PA-BA-26-1 PA-BA-92 PA-BA-40 PA-BA-127 PA-PP-XPP PA-BA-23-A PA-BA-23-B | 30 000 Max 10 000 30 000 30 000 30 000 / NEW 30 000 NEW Max 10 000 NEW 30 000 | 36 days 34 days 34 days 40 days 40 days 10 days 33 days 33 days | 219 245 170 295 346 299 76 352 228 |













| Mycology - See p.8 "Seed health" | | | | |
|--|----------------|------|----------|--------|
| | | Size | Duration | Price |
| Watermelon | | | | |
| Stagonosporopsis cucurbitacearum (Didymella bryoniae), Fusarium oxysporum, Fusasolani, Alternaria cucumerina, Gloeosporium orbiculare (Colletotrichum orbiculare), Fusarium (other sections), Phomopsis vexans, Botrytis sp., Cladosporium sp. | arium | | | |
| Agar method with superficial disinfection. UNTREATED seeds only. | PA-ES-PASD | 400 | 19 days | 110.00 |
| Agar method without superficial disinfection. Treated seeds only. | PA-ES-PAS | 400 | 19 days | 106.00 |
| Capsicum | | | | |
| Phytophthora capsici | | | | |
| Seed wash method. UNTREATED seeds only. | PA-MI-PIM | 500 | 15 days | 102.00 |
| Capsicum, Pepper | | | | |
| Colletotrichum truncatum (Colletotrichum capsici), Fusarium oxysporum, Fusarium (a Colletotrichum coccodes, Sclerotinia sclerotiorum, Botrytis sp., Verticillium sp. | all sections), | | | |
| Agar method. | PA-ES-POIV | 400 | 19 days | 106.00 |
| Leek | | | | |
| Alternaria porri, Botrytis allii and/or Botrytis aclada, Sclerotinia minor, Fusarium mo | niliforme, | | | |
| Fusarium oxysporum, Fusarium (other sections), Botrytis sp., Stemphylium sp. | DA EC DOD | 400 | 10 4 | 100.00 |
| Agar method. Pea | PA-ES-POR | 400 | 19 days | 106.00 |
| | alla | | | |
| Didymella pisi (Ascochyta pisi), Didymella pinodes (Mycosphaerella pinodes), Didym pinodella (Phoma pinodella), Stemphylium botryosum, Fusarium (other sections), Bot Sclerotinia sp., Phoma sp. | | | | |
| Agar method with superficial disinfection. UNTREATED seeds only. | PA-ES-POID | 400 | 19 days | 110.00 |
| Agar method without superficial disinfection. Treated seeds only. | PA-ES-POI | 400 | 19 days | 106.00 |
| Peronospora viciae (Peronospora pisi) (downy mildew) Seed wash method. UNTREATED seeds only. | PA-MI-POI | 500 | 15 days | 102.00 |
| Didymella pisi (Ascochyta pisi) | | | | |
| Agar method (ISTA 7-005). | PA-ANT-POI | 400 | 19 days | 110.00 |
| Chickpea | | | | |
| Ascochyta rabiei (Phoma rabiei), Botrytis cinerea, Fusarium oxysporum, Fusarium so. Fusarium (other sections) | lani, | | | |
| Agar method with superficial disinfection. UNTREATED seeds only. | PA-ES-POCD | 400 | 19 days | 110.00 |
| Agar method without superficial disinfection. Treated seeds only. | PA-ES-POC | 400 | 19 days | 106.00 |
| Radish | | | | |
| Hyaloperonospora parasitica (Peronospora parasitica) (downy mildew) | | | | |
| Seed wash method. UNTREATED seeds only. | PA-MI-RAD | 500 | 15 days | 102.00 |
| Grow-out method (viability testing). | PA-MIRADGO | 400 | 42 days | 129.00 |
| Rocket | | | | |
| Hyaloperonospora parasitica Seed wash method. UNTREATED seeds only. | PA-MI-ROQL | 500 | 15 days | 102.00 |
| Tomato | PA-IVII-ROQL | 300 | 13 days | 102.00 |
| Alternaria solani, Fusarium oxysporum, Fusarium solani, Colletotrichum coccodes, Botrytis cinerea, Fusarium (all sections), Didymella sp., Verticillium sp., | | | | |
| Stemphylium sp., Rhizoctonia sp., Sclerotinia sp. | | | | |
| Agar method. | PA-ES-TOM | 400 | 19 days | 106.00 |
| N . 1 | | | | |
| Nematology | | | | |
| | | Size | Duration | Price |
| Carrot | | | | |
| Ditylenchus dipsaci Filtration (COFRAC - Anses MOA013 parts A and B). UNTREATED seeds only. Test carried out on the whole submitted sample. If the supplied quantity is too important, a new sample will be requested. | PA-NE-CAR | 70 g | 16 days | 76.00 |
| Onion | | | | |
| Ditylenchus dipsaci | | | | |
| Filtration (COFRAC - Anses MOA013 parts A and B). UNTREATED seeds only. Test carried out on the whole submitted sample. If the supplied quantity is too important, | PA-NE-OIG | 70 g | 16 days | 76.00 |
| a new sample will be requested. | | | | |



Vegetables •—



| Virology - Uncoated seeds only | | Size | Duration | Price |
|--|------------|-------|----------|--------|
| Eggplant, Capsicum, Pepper, Tomato | | | | |
| Tomato mosaic virus (ToMV) and/or Tobacco mosaic virus (TMV) | | | | |
| ELISA. | PA-VI-19 | 3 000 | 16 days | 185.00 |
| Tobacco mild green mosaic virus (TMGMV) ELISA. | PA-VI-94-1 | 1 000 | 16 days | 136.00 |
| ELI3A. | PA-VI-94-1 | 3 000 | 16 days | 147.00 |
| Tomato brown rugose fruit virus (ToBRFV)40 | | | | |
| Supplement fee. In addition analysis with ISHI primers on request. | PA-VI-ISHI | 1 | 10 days | 137.00 |
| Supplement fee. Confirmation by PCR of ELISA positive subsamples. | PA-VI-PCR | 1 | / | 405.00 |
| Arabis mosaic virus (ArMV) | | | | |
| ELISA. | PA-VI-33-1 | 3 000 | 16 days | 220.00 |
| Tomato bushy stunt virus (TBSV) | DA VI 47 | 2 000 | 16 days | 222.00 |
| Delargenium zenate enet virus (DTSVI) | PA-VI-47 | 3 000 | 16 days | 232.00 |
| Pelargonium zonate spot virus (PZSV) ELISA. | PA-VI-46 | 3 000 | 16 days | 246.00 |
| Pepper veinal mottle virus (PVMV) | | | | |
| ELISA. | PA-VI-86 | / | / | 235.00 |
| Pepino mosaic virus (PepMV) | | | | |
| ELISA and confirmation of positives and indeterminates by RT-PCR. (Internal method | PA-VI-15 | 1 000 | 16 days | 168.00 |
| derived from Anses MOA 008 – MOA 026). | PA-VI-16 | 3 000 | 16 days | 195.00 |
| _ | PA-VI-17 | 5 000 | 16 days | 335.00 |
| Carrot | | | | |
| Alfalfa mosaic (AMV) | | | | |
| ELISA. | PA-VI-71 | 2 000 | 16 days | 162.00 |
| Arabis mosaic virus (ArMV) | | | | |
| ELISA. | PA-VI-33-1 | 3 000 | 16 days | 220.00 |
| Cucumber leaf spot carmovirus (CLSV) ELISA. | PA-VI-35-1 | 3 000 | 16 days | 220.00 |
| Tomato ringspot virus (ToRSV) ⁴⁰ | FA-VI-33-1 | 3 000 | 16 days | 220.00 |
| ELISA. | PA-VI-38-1 | 3 000 | 16 days | 162.00 |
| Celery | | | | |
| Strawberry latent ringspot virus (SLRSV) | | | | |
| ELISA. | PA-VI-36 | / | / | 250.00 |
| Cucumis sp. | | | | |
| Cucumber leaf spot carmovirus (CLSV) ELISA. | PA-VI-35-1 | 3 000 | 16 days | 220.00 |
| Cucurbits - Detection of 1 pathogen | 14 (133 1 | 3 000 | 10 days | 220.00 |
| Cucumber mosaic virus (CMV) | | | | |
| ELISA. | PA-VI-56 | 2 000 | 16 days | 243.00 |
| Arabis mosaic virus (ArMV) | | | | |
| ELISA. | PA-VI-33 | 2 000 | 16 days | 247.00 |
| Cucumber leaf spot carmovirus (CLSV) | DA 1// 25 | 2.000 | 16 - | 247.00 |
| ELISA. | PA-VI-35 | 2 000 | 16 days | 247.00 |
| Tobacco ringspot virus (TRSV) ⁴⁰ ELISA. | PA-VI-39 | 2 000 | 16 days | 258.00 |
| Tomato black ring virus (TBRV) | 17, 17, 33 | 2 000 | 10 0045 | |
| ELISA. | PA-VI-37 | 2 000 | 16 days | 205.00 |
| Tomato ringspot virus (ToRSV) ⁴⁰ | | | | |
| ELISA. | PA-VI-38 | 2 000 | 16 days | 247.00 |
| Zucchini yellow mosaic virus (ZYMV) | | | | |
| ELISA. | PA-VI-40 | 2 000 | 16 days | 255.00 |
| Squash leaf curl virus (SLCV) | 54.1/2.== | , | , | 250.55 |
| ELISA. | PA-VI-77 | / | | 250.00 |
| Cucumber green mottle mosaic virus (CGMMV) ELISA (ISTA 7-026). | PA-VI-01-1 | 2 000 | 16 days | 185.00 |
| LLION (ISTN 1-020). | PA-VI-01-1 | 2 000 | TO UdyS | 103.00 |



| Virology - Uncoated seeds only | | Size | Duration | Price |
|---|---|----------|----------|--------|
| Lettuce | | 3120 | Daration | |
| Tomato ringspot virus (ToRSV) ⁴⁰ | | | | |
| ELISA. | PA-VI-38-1 | 3 000 | 16 days | 162.0 |
| Strawberry latent ringspot virus (SLRSV) | | | | |
| ELISA. | PA-VI-36 | / | | 250.0 |
| Capsicum, Pepper, Tomato | | | | |
| Tomato brown rugose fruit virus (ToBRFV) ⁴⁰ RT-PCR (COFRAC - method ANSES/LSV/MA066, primers and probe of Menzel and Winter). UNCOATED seeds olny. | PA-VI-93-7 | 1 000 | 10 days | 146.0 |
| <u> </u> | PA-VI-93-6 | 3 000 | 10 days | 196.0 |
| RT-PCR (COFRAC - method ANSES/LSV/MA066, primers and probe of Menzel and Winter). | PA-VI-9310 NEV | V 20 000 | 10 days | 2376.0 |
| Tomato brown rugose fruit virus (ToBRFV) ⁴⁰ and Tomato mottle mosaic virus (ToMMV RT-PCR (COFRAC - method ANSES/LSV/MA066, primers and probe of Menzel and Winter for ToBRFV). | PA-VI-99-1 NEV | V 1 000 | 10 days | 157.0 |
| | PA-VI-99 NEV | V 3 000 | 10 days | 238.0 |
| | PA-VI-99-4 NEV | V 20 000 | 10 days | 2700.0 |
| Pea early browning virus (PEBV) ELISA (ISTA 7-024). | PA-VI-31 | 2 000 | 16 days | 205.0 |
| Pea enation mosaic virus (PEMV) | DA 1/1 57 | 2 000 | 16 4 | 240.0 |
| ELISA. | PA-VI-57 | 2 000 | 16 days | 249.0 |
| Pea seed borne mosaic virus (PSbMV) ELISA (ISTA 7-024). | PA-VI-11 | 2 000 | 16 days | 173.0 |
| Bean yellow mosaic virus (BYMV) | .,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,, | 2 000 | | 17510 |
| ELISA. | PA-VI-60 | / | / | 271.0 |
| Bean leaf roll virus (BLRV) | | | | |
| ELISA. | PA-VI-67 | / | / | 247.0 |
| Southern bean mosaic virus (SBMV) ELISA. | PA-VI-88 | / | / | 247.0 |
| Tomato | | | | |
| Pepino mosaic virus (PepMV) ELISA (COFRAC - method Anses MOA 026) and confirmation of positives and ndeterminates according by RT-PCR (Internal method). | PA-VI-15CO | 1 000 | 16 days | 165.0 |
| | PA-VI-16CO | 3 000 | 16 days | 242.0 |
| | PA-VI-17CO | 5 000 | 16 days | 398.0 |


EVALUATION OF VARIETIES

| Varietal resistance | | | | |
|---|------------|------|----------|--------|
| | | Size | Duration | Price |
| Eggplant | | | | |
| Verticillium dahliae | | | | |
| GEVES protocol. | PA-R-AUB-1 | 45 | / | 186.00 |
| Cabbage | | | | |
| Fusarium oxysporum f. sp. conglutinans race 1 | | | | |
| Official protocol. | PA-R-CHO | 45 | / | 330.00 |
| Plasmodiophora brassicae | | | | |
| GEVES protocol. | PA-R-CHO-1 | 45 | / | 242.00 |
| Cucumber | | | | |
| CMV (Cucurbit mosaic virus) | | | | |
| Official protocol. | PA-R-CON | 45 | / | 150.00 |
| CGMMV (Cucumber green mottle mosaic virus) | | | | |
| GEVES protocol. | PA-R-CON-1 | 45 | | 160.00 |
| ZYMV (Zucchini yellow mosaic virus) | | | | |
| Official protocol. | PA-R-CON-2 | 45 | / | 160.00 |
| | | | | |

Different prices outside test periods. Contact SNES for information on the periods according to the species.

| Varietal resistance | | Size | Duration | Price |
|--|-------------------|------|-----------|-----------|
| Cucumber | | 3126 | Duration | Price |
| WMV (Watermelon mosaic virus) | | | | |
| Official protocol. | PA-R-CON-3 | 45 | / | 160.00 |
| Podosphaera xanthii race 1 | | | | |
| Official protocol. | PA-R-CON-4 | 45 | | 313.00 |
| Squash | | | | |
| ZYMV (Zucchini yellow mosaic virus) | | 4.5 | , | 460.00 |
| Official protocol. | PA-R-COU-2 | 45 | / | 160.00 |
| WMV (Watermelon mosaic virus) Official protocol. | PA-R-COU-3 | 45 | / | 159.00 |
| Podosphaera xanthii race 1 | | | <u> </u> | |
| Official protocol. | PA-R-COU-4 | 45 | / | 304.00 |
| Strawberry | | | | |
| Phytophthora cactorum | | | | |
| Official protocol. | PA-R-FRA-1 | | Conta | act SNES |
| Bean | | | | |
| BCMNV (Bean common mosaic necrotic virus) | DA D 114D 4 | 20 | , | 427.00 |
| Official protocol. | PA-R-HAR-1 | 30 | / | 127.00 |
| Colletotrichum lindemuthianum race 6 or race Kappa (anthracnose) | PA-R-HAR-COL | 30 | / | 141.00 |
| Pseudomonas savastanoi pv. phaseolicola race 6 (halo blight) | r A-N-HAN-COL | | | 141.00 |
| Official protocol. | PA-R-HAR-3 | 30 | / | 171.00 |
| Xanthomonas axonopodis pv. phaseoli | - | | | |
| Official protocol. | PA-R-HAR-4 | | Conta | act SNES |
| Lettuce | | | | |
| Bremia lactucae races Bl: 1EU / Bl: 2EU / Bl: 3EU / Bl: 4EU / Bl: 5EU / Bl: 6EU / Bl: 7EU Bl: 10EU / Bl: 12 à 15EU / Bl: 17EU / Bl: 18EU / Bl: 20EU / Bl: 22 à 25EU / Bl: 28EU / Bl: 32EU / Bl: 34EU / S1 / SF1 ou IL4 | / | | | |
| Official protocol. | PA-R-LAI-BRE | | Conta | act SNES |
| Bremia lactucae official races for CTPS BI: 16EU / BI: 21EU / BI: 26EU / BI: 27EU / BI: 29EU / BI: 31EU / BI: 33EU / BI: 35EU ou BI: 36EU | | | | |
| Official protocol. | PA-R-LAI-BRE1 | 45 | / | 66.00 |
| Bremia lactucae new races Bl: 37EU / Bl: 38EU / Bl: 39 EU and Bl: 40EU | DA D LALDDES ALEM | 45 | , | 66.00 |
| Official protocol. | PA-R-LAI-BRE2 NEW | 45 | / | 66.00 |
| Bremia lactucae Late stage resistance. | PA-R-LAI29 | | Cont | act SNES |
| Identification of the race. | PA-R-IDBRE | | | act SNES |
| LMV (<i>Lettuce mosaic virus</i>) pathotype II (LMV-0) or pathotype III (LMV-9) | | | | 301 0.120 |
| Official protocol. | PA-R-LAI-LMV | | Conta | act SNES |
| LMV (Lettuce mosaic virus) | | | | |
| Detection of markers linked to resistance genes. Gene mo1. Resistance to the Lettuce virus. | BI-D-GENR | | Contact B | ioGEVES |
| Fusarium oxysporum f. sp. lactucae race 1 | | | | |
| Official protocol. | PA-R-LAI30 | 45 | / | 179.00 |
| Fusarium oxysporum f.sp. lactucae race 4 Official protocol. | PA-R-LAI41 NEW | 45 | / | 195.00 |
| Fusarium oxysporum f. sp. lactucae | | | | |
| Identification of the race. | PA-R-IDFUS | | Conta | act SNES |
| Nasonovia ribisnigri race 0 | | | | |
| Official protocol. | PA-R-LAI35 | 45 | / | 172.00 |
| Corn salad | | | | |
| Peronospora valerianellae race Pv: 1 or Pv: 2 Official protocol. | PA-R-MAC-PV | | Conta | act SNES |
| Melon | | | | |
| Fusarium oxysporum f. sp. melonis races Fom: 0 / Fom: 1 / Fom: 2 or Fom: 1.2 | DA D MEL EUS | ΔF | , | 175 00 |
| Official protocol. | PA-R-MEL-FUS | 45 | / | 175.00 |

Different prices outside test periods. Contact SNES for information on the periods according to the species.



Different prices outside test periods. Contact SNES for information on the periods according to the species.

Vegetables •—

| Varietal resistance | | | | |
|---|--------------------|------|-----------|----------|
| Tomato | | Size | Duration | Price |
| ToMV: 0, 1, 2 (<i>Tomato mosaic virus</i>) races ToMV: 0 / ToMV: 1 ou ToMV: 2 | | | | |
| Official protocol. | PA-R-TOM-TMV | 45 | / | 165.00 |
| TMV: 0 (Tobacco mosaic virus) | | | | |
| Detection of markers linked to resistance genes. Gene Tm1. Resistance to the Tomatovirus. | BI-D-GENR | | Contact B | ioGEVES |
| Detection of markers linked to resistance genes. Tm2 and Tm2 ² genes. Resistance to the Tomatovirus. | BI-D-GENR | | Contact B | ioGEVES |
| TSWV: 0 (<i>Tomato spotted wilt virus</i>) race 0 Official protocol. | PA-R-TOM10 | 60 | / | 165.00 |
| Meloidogyne incognita Official protocol. | PA-R-TOM14 | 45 | / | 115.00 |
| Pseudomonas syringae pv. tomato Official protocol. | PA-R-TOM15 | 45 | / | 91.00 |
| Pseudopyrenochaeta lycopersici Official protocol. | PA-R-TOM16 | 60 | , | 398.00 |
| Passalora fulva | FA-R-TOWID | 00 | | 336.00 |
| Identification of the race. | PA-ID-PF | | Conta | act SNES |
| Tomato rootstock | | | | |
| Verticillium dahliae Official protocol. | PA-R-TPG-1 | 90 | / | 177.00 |
| Fusarium oxysporum f. sp. lycopersici races Fol: 0 / Fol: 1 or Fol: 2 Official protocol. | PA-R-TPG-FUS | 90 | / | 185.00 |
| Passalora fulva races Pf: 0 or Pf: E Official protocol. | PA-R-TPG-PF | 90 | / | 177.00 |
| Fusarium oxysporum radicis f. sp. lycopersici Official protocol. | PA-R-TPG-7 | 90 | | 177.00 |
| Stemphylium spp. Official protocol. | PA-R-TPG-8 | 90 | | 177.00 |
| ToMV: 0, 1, 2 (Tomato mosaic virus) races ToMV: 0 / ToMV: 1 ou ToMV: 2 | | | , | |
| Official protocol. | PA-R-TPG-TMV | 90 | / | 170.00 |
| TSWV: 0 (Tomato spotted wilt virus) race 0 Official protocol. | PA-R-TPG10 | 90 | / | 170.00 |
| Meloidogyne incognita | DA D TDC14 | 00 | , | 117.00 |
| Official protocol. Pseudomonas syringae pv. tomato | PA-R-TPG14 | 90 | | 117.00 |
| Official protocol. | PA-R-TPG15 | 90 | / | 95.00 |
| Pseudopyrenochaeta lycopersici Official protocol. | PA-R-TPG16 | 90 | / | 393.00 |
| Different prices outside test periods. Contact SNES for information on the periods according | | | , | |
| Genotyping by molecular biology | | | | |
| Cabbage, Strawberry, Lettuce, Pea, Radish | | Size | Duration | Price |
| Varietal identity control - SSR. | BI-G-BM-SSR-CID-1 | | Contact B | ioGEVES |
| Varietal purity analysis - SSR - 90 seeds. | BI-G-BM-SSR-PUR-90 | | Contact B | ioGEVES |
| Technological quality: biochemicals tests | | | | |
| Cabbage, Radish, Other Brassicaceae | | Size | Duration | Price |
| Glucosinolate content (HPLC). | BI-B-HPLC-GLU | | Contact B | ioGEVES |
| Fatty acid composition (CPG method). | BI-B-CPG-AG | | Contact B | ioGEVES |
| Field Bean, Pea Protein content (NIRS). | BI-B-NIRS-P | | Contact B | ioGFVFS |
| Capsicum/Pepper | טו ט־וווווק-ר | | Contact B | .552423 |
| Capsaicin and dihydrocapsaicin content (capsaicinoids) (HPLC). | BI-B-HPLC-CAP | | Contact B | ioGEVES |



Technological quality: biochemicals tests

Size Duration Price

Pea

Contact BioGEVES BI-B-SPEC-FAT Antitrypsic factors (assay by spectrophotometry).

| Field tests by SEV | | |
|---|-----------------|---------|
| | | Price |
| DUS testing - Cucumber, Lettuce, Melon, Pepper, Tomato Cycle 1. | SEV-DHS-POTMAJ1 | 1990.00 |
| DUS testing - Cucumber, Lettuce, Melon, Pepper, Tomato Cycle 2. | SEV-DHS-POTMAJ2 | 1880.00 |
| DUS testing - Other vegetables species Cycle 1. | SEV-DHS-POTMIN1 | 1325.00 |
| DUS testing - Other vegetables species Cycle 2. | SEV-DHS-POTMIN2 | 1240.00 |

PUBLICATIONS (Contact SNES)

Method sheet

| Vigour testing - Conductivity - Pea. | VIG-2-M |
|---|------------|
| Germination analysis technical sheet | |
| Ealuation of Carrot seedlings. | GE-T-CAR |
| Evaluation of Cabbage seedlings. | GE-T-CHOU |
| Evaluation of Bean seedlings. | GE-T-HAR |
| Evaluation of Lettuce seedlings. | GE-T-LAI |
| Evaluation of Onion seedlings. | GE-T-OIG |
| Evaluation of Pea seedlings. | GE-T-POI |
| Evaluation of Radish seedlings. | GE-T-RAD |
| Evaluation of Tomato seedlings. | GE-T-TOM |
| Technical sheet for analysis of specific purity and counting of all other seeds | |
| Pisum sativum, Vicia faba. | AP-C-8 |
| Cicer arietinum. | AP-C-12 |
| Allium sp. (Allium cepa, Allium porrum, Allium schoenoprasum). | AP-C-13 |
| Solanaceae. (Solanum lycopersicum, Solanum melongena, Capsicum annuum). | AP-C-14 |
| Daucus carota, Petroselinum sp. | AP-C-15 |
| Cucurbitaceae. (Curcurbita spp., Cucumis spp., Citrullus Ianatus). | AP-C-16 |
| Identification data sheet of seeds and other impurities | |
| Asteraceae (Anthemis arvensis, Glebionis segetum, Chicorium sp., Tripleurospermum inodorum, Helminthotheca echioïdes, | AP-A-06 |
| Lapsana communis, Lactuca sativa, Sonchus spp., Cirsium arvense, Cirsium vulgare, Centaurea cyanus). | |
| Collection of seeds | |
| Weed's identification for <i>Pisum sativum</i> and <i>Vicia faba</i> analysis. | APCS-PIS-S |
| Weed's identification for Vegetables analysis. | APCS-VEG |
| | |

Ornamental and Fruit crops

SEED QUALITY Physical quality Duration Size Price Thousand-seed weight (on purity test performed by SNES) Thousand-seed weight on pure seeds. MMS-01 33.00 **Purity analysis test** ISTA weight Purity - Fruit crops, Ornamentals. PU-IS-18 33.20 Percentage of a specific type of other seeds. Specify the search to be performed. PU-CONS1 9.00 PU-CONS2 9.00 Percentage of a specific type of inert materials. Specify the search to be performed. PU-LB-SUP Contact SNES Supplement for purity analysis if received as raw seeds. Counting of all other seeds SP-IS-17 ISTA weight 138.00 Full counting - Fruit crops, Ornamentals. Counting of other seeds on purity weight. Indication of the number of other seeds in the specific PU-SP-01 13.40 purity test. Limited counting of all other seeds Searching of 1 to 4 species (except for Orobanchaceae). Indicate the name of the species to be SP-LI-01 ISTA weight 64.00 Searching of 5 to 8 species (except for Orobanchaceae). Indicate the name of the species to be SP-LI-02 102.00 ISTA weight searched. SP-LI-20 Contact SNES Searching of more than 8 species (except for Orobanchaceae). Indicate the name of the species to be searched. Moisture content - Provide seeds in watertight bags from which as much air as possible has been extracted Oven method. TE-SN-01 ISTA weight 20.70 Identification of individual seeds Visual identification by species. ID-IS-01 34.60 Physiological quality Size Duration Price Germination test on 400 seeds Trees, Shrubs, Flowers. GE-FG-20-4 1 250 76.00 Germination test on 200 seeds Trees, Shrubs, Flowers. GE-FG-20-2 500 60.00 Germination tests on bulbs and bulblets 1 250 151.00 On 400 seeds. **GE-BULB-4** GE-BULB-2 122.00 On 200 seeds. 500 Tetrazolium viability test on 400 seeds - For results within a week, reception of seeds on Tuesday at the latest. Oak, Dogwood, Olive, Hazelnut, Walnut. GE-TZ-3-4 500 219.00 Hornbeam, Maple, Ash, Stone fruits, Beech, Lavender, Rosemary. **GE-TZ-2-4** 500 185.00 GE-TZ-1-4 500 174.00 Amelanchier, Conifers, Ligustrum, Mahonia, Apple, Pear, Sorbier, . Tetrazolium viability test on 200 seeds - For results within a week, reception of seeds on Tuesday at the latest. GE-TZ-3-2 300 150.00 Oak, Dogwood, Olive, Hazelnut, Walnut. Hornbeam, Maple, Ash, Stone fruits, Beech, Lavender, Rosemary. 300 127.00 GE-TZ-2-2 Amelanchier, Conifers, Ligustrum, Mahonia, Apple, Pear, Sorbier, . 300 116.00 GE-TZ-1-2 Tetrazolium viability test on 100 seeds - For results within a week, reception of seeds on Tuesday at the latest. 200 116.00 Oak, Dogwood, Olive, Hazelnut, Walnut. GF-TZ-3-1 Hornbeam, Maple, Ash, Stone fruits, Beech, Lavender, Rosemary. GE-TZ-2-1 200 93.00 Amelanchier, Conifers, Ligustrum, Mahonia, Apple, Pear, Sorbier, . GE-TZ-1-1 200 81.00 Verification of species Verification of species after germination test. **GE-ENR** 9.40

Ornamental and Fruit crops

| Nematology Size Duration Price Bulbs (COFRAC), bulblets, corms, rhizomes, tubers Ditylenchus dipsaci Filtration (Anses MOA013 parts A and B). UNTREATED seeds only. Test carried out on the whole submitted sample. If the supplied quantity is too important, a new sample will be requested. Virology - Uncoated seeds only Size Duration Price Cyclamen Tomato spotted wilt virus (TSWV) ELISA. PA-VI-49 / / 262.00 |
|---|
| Size Duration Price Bulbs (COFRAC), bulblets, corms, rhizomes, tubers Ditylenchus dipsaci Filtration (Anses MOA013 parts A and B). UNTREATED seeds only. Test carried out on the whole submitted sample. If the supplied quantity is too important, a new sample will be requested. Virology - Uncoated seeds only Cyclamen Tomato spotted wilt virus (TSWV) ELISA. PA-VI-49 PA-VI-49 PA-VI-49 PA-VI-49 PA-VI-49 PA-VI-49 PA-VI-49 PA-VI-49 PA-VI-49 |
| Bulbs (COFRAC), bulblets, corms, rhizomes, tubers Ditylenchus dipsaci Filtration (Anses MOA013 parts A and B). UNTREATED seeds only. Test carried out on the whole submitted sample. If the supplied quantity is too important, a new sample will be requested. Virology - Uncoated seeds only Size Duration Price Cyclamen Tomato spotted wilt virus (TSWV) ELISA. PA-VI-49 / / 262.00 |
| Ditylenchus dipsaci Filtration (Anses MOA013 parts A and B). UNTREATED seeds only. Test carried out on the whole submitted sample. If the supplied quantity is too important, a new sample will be requested. Virology - Uncoated seeds only Size Duration Price Cyclamen Tomato spotted wilt virus (TSWV) ELISA. PA-VI-49 / / 262.00 |
| Filtration (Anses MOA013 parts A and B). UNTREATED seeds only. Test carried out on the whole submitted sample. If the supplied quantity is too important, a new sample will be requested. Virology - Uncoated seeds only Size Duration Price Cyclamen Tomato spotted wilt virus (TSWV) ELISA. PA-VI-49 / / 262.00 |
| Test carried out on the whole submitted sample. If the supplied quantity is too important, a new sample will be requested. Virology - Uncoated seeds only Size Duration Price Cyclamen Tomato spotted wilt virus (TSWV) ELISA. PA-VI-49 / 262.00 EVALUATION OF VARIETIES |
| Virology - Uncoated seeds only Size Duration Price Cyclamen Tomato spotted wilt virus (TSWV) ELISA. PA-VI-49 / / 262.00 EVALUATION OF VARIETIES |
| Size Duration Price Cyclamen Tomato spotted wilt virus (TSWV) ELISA. PA-VI-49 / / 262.00 EVALUATION OF VARIETIES |
| Size Duration Price Cyclamen Tomato spotted wilt virus (TSWV) ELISA. PA-VI-49 / / 262.00 EVALUATION OF VARIETIES |
| Cyclamen Tomato spotted wilt virus (TSWV) ELISA. PA-VI-49 / / 262.00 EVALUATION OF VARIETIES |
| Tomato spotted wilt virus (TSWV) ELISA. PA-VI-49 / / 262.00 EVALUATION OF VARIETIES |
| EVALUATION OF VARIETIES PA-VI-49 / / 262.00 |
| EVALUATION OF VARIETIES |
| |
| |
| |
| Genotyping by molecular biology |
| Size Duration Price |
| Apricot, Cherry tree, Hydrangea, Kiwi, Hazel tree, Walnut tree, Palm, Peach, |
| Poplar, Apple Tree, Pear Tree, Plum tree, Willow |
| Varietal identity control - SSR. BI-G-BM-SSR-CID-1 Contact BioGEVES |
| Quince |
| Varietal identity control. BI-G-BM-SSR-CID-9 NEW Contact BioGEVES |
| Palm |
| Varietal identity control for export (True-to-type nature). BI-G-BM-SSR-CID-6 Contact BioGEVES |
| Varietal identity control for production (True-to-type nature). BI-G-BM-SSR-CID-7 Contact BioGEVES |
| Poplar Varietal identity control among french cultivars. BI-G-BM-SSR-CID-8 Contact BioGEVES |
| varietal facility control allong french calcivals. |
| Bud sample for genotyping |
| Price |
| Cost of sampling for 1 INRAE site and 1 applicant/breeder. SEV-ECHF-FOR 376.00 |
| Cost for 1 sampled variety. SEV-ECHF-VAR 41.00 |
| Packaging by INRAE examiner for 1 site and for 1 to 5 varieties. SEV-ECHF-COND5 150.00 |
| Packaging by INRAE examiner for 1 site and for 6 to 10 varieties. SEV-ECHF-COND10 300.00 |
| Packaging by INRAE examiner for 1 site and for 11 to 50 varieties. SEV-ECHF-COND50 623.00 |
| Cost of sending for 1 site (possible to pick the samples directly on the site). SEV-ECHF-ENV 125.00 |
| |
| Field tests by SEV |
| Price Price Price |
| DUS testing - Fruit trees and rootstock - New variety, installation year. SEV-DHS-FRU1 1100.00 |
| DUS testing - Fruit trees and rootstock - New variety, following years. SEV-DHS-FRU2 2200.00 |
| DUS testing - Ornementals species. SEV-DHS-ORN 2800.00 |
| DUS testing - Vine - New variety, installation year. SEV-DHS-VIG1 1100.00 |
| DUS testing - Vine - New variety, following years. SEV-DHS-VIG2 2200.00 |
| |

PUBLICATIONS (Contact SNES)

Identification data sheet of seeds and other impurities

Lathyrus spp. (Lathyrus sylvestris, Lathyrus latifolius, Lathyrus hirsutus, Lathyrus tuberosus, Lathyrus odoratus, Lathyrus aphaca, Lathyrus pratensis, Lathyrus sativus, Lathyrus cicera).

AP-A-05

Aromatic medicinal

| Physical quality | | | | |
|--|-------------------------------|----------------------------|-------------------------|--------------------------------------|
| | | Size | Duration | Price |
| Thousand-seed weight (on purity test performed by SNES) | | | | |
| Thousand-seed weight on pure seeds. | MMS-01 | /_ | /_ | 33.0 |
| Purity analysis test | | | | |
| Purity - Aromatic, Medicinal. | PU-IS-18 | ISTA weight | /_ | 33.20 |
| Percentage of a specific type of other seeds. Specify the search to be performed. | PU-CONS1 | / | /_ | 9.00 |
| Percentage of a specific type of inert materials. Specify the search to be performed. | PU-CONS2 | / | /_ | 9.00 |
| Supplement for purity analysis if received as raw seeds. | PU-LB-SUP | | Conta | act SNES |
| Counting of all other seeds | | | , | |
| Full counting - Aromatic, Medicinal. | SP-IS-17 | ISTA weight | | 138.00 |
| Counting of other seeds on purity weight. Indication of the number of other seeds in the specific purity test. | PU-SP-01 | / | / | 13.4 |
| Limited counting of all other seeds | | | | |
| Searching of 1 to 4 species (except for <i>Orobanchaceae</i>). Indicate the name of the species to be searched. | SP-LI-01 | ISTA weight | / | 64.00 |
| Searching of 5 to 8 species (except for <i>Orobanchaceae</i>). Indicate the name of the species to be searched. | SP-LI-02 | ISTA weight | / | 102.00 |
| Searching of more than 8 species (except for <i>Orobanchaceae</i>). Indicate the name of the species to be searched. | SP-LI-20 | | Conta | act SNES |
| Moisture content - Provide seeds in watertight bags from which as much air as | | | | |
| possible has been extracted Oven method. | TE-SN-01 | ISTA weight | | 20.70 |
| Identification of individual seeds | | | | |
| Visual identification by species. | ID-IS-01 | / | / | 34.60 |
| Germination test on 400 seeds | | Size | Duration | Price |
| Aromatics, Medicinals. | GE-FG-18-4 | 1 250 | / | 65.00 |
| Germination test on 200 seeds | | | | |
| Aromatics, Medicinals. | GE-FG-18-2 | 500 | | 52.00 |
| Bacteriology - Uncoated seeds only | | | | |
| | | Size | Duration | Price |
| Dill Coriander Parsley - Detection of 1 nathogen | | | | |
| Dill, Coriander, Parsley - Detection of 1 pathogen | | | | |
| Pseudomonas viridiflava | PA-RA-10 <i>1</i> | 30,000 | 26 days | 31/1 0 |
| Pseudomonas viridiflava Agar method + PCR in case of suspect colonies. | PA-BA-104 | 30 000 | 26 days | 314.00 |
| Pseudomonas viridiflava Agar method + PCR in case of suspect colonies. Pseudomonas syringae pv. apii | | | | 314.00 |
| Pseudomonas viridiflava Agar method + PCR in case of suspect colonies. Pseudomonas syringae pv. apii Agar method + PCR in case of suspect colonies. | PA-BA-104 PA-BA-106 | 30 000 30 000 | 26 days | 314.00 278.00 |
| Pseudomonas viridiflava Agar method + PCR in case of suspect colonies. Pseudomonas syringae pv. apii Agar method + PCR in case of suspect colonies. Pseudomonas syringae pv. coriandricola | PA-BA-106 | 30 000 | 36 days | 278.00 |
| Pseudomonas viridiflava Agar method + PCR in case of suspect colonies. Pseudomonas syringae pv. apii Agar method + PCR in case of suspect colonies. Pseudomonas syringae pv. coriandricola Agar method + PCR in case of suspect colonies. | | | | 278.00 |
| Pseudomonas viridiflava Agar method + PCR in case of suspect colonies. Pseudomonas syringae pv. apii Agar method + PCR in case of suspect colonies. Pseudomonas syringae pv. coriandricola Agar method + PCR in case of suspect colonies. Dill, Coriander, Parsley - Detection of 2 pathogens | PA-BA-106 | 30 000 | 36 days | 278.0 |
| Pseudomonas viridiflava Agar method + PCR in case of suspect colonies. Pseudomonas syringae pv. apii Agar method + PCR in case of suspect colonies. Pseudomonas syringae pv. coriandricola Agar method + PCR in case of suspect colonies. Dill, Coriander, Parsley - Detection of 2 pathogens Pseudomonas syringae pv. apii + Pseudomonas syringae pv. coriandricola | PA-BA-106 PA-BA-107 | 30 000 30 000 | 36 days 26 days | 278.00 |
| Pseudomonas viridiflava Agar method + PCR in case of suspect colonies. Pseudomonas syringae pv. apii Agar method + PCR in case of suspect colonies. Pseudomonas syringae pv. coriandricola Agar method + PCR in case of suspect colonies. Dill, Coriander, Parsley - Detection of 2 pathogens Pseudomonas syringae pv. apii + Pseudomonas syringae pv. coriandricola Agar method + PCR in case of suspect colonies. | PA-BA-106 | 30 000 | 36 days | 278.00 |
| Pseudomonas viridiflava Agar method + PCR in case of suspect colonies. Pseudomonas syringae pv. apii Agar method + PCR in case of suspect colonies. Pseudomonas syringae pv. coriandricola Agar method + PCR in case of suspect colonies. Dill, Coriander, Parsley - Detection of 2 pathogens Pseudomonas syringae pv. apii + Pseudomonas syringae pv. coriandricola Agar method + PCR in case of suspect colonies. Pseudomonas syringae pv. apii + Pseudomonas viridiflava | PA-BA-106 PA-BA-107 | 30 000 30 000 | 36 days 26 days | 278.00 289.00 359.00 |
| Pseudomonas viridiflava Agar method + PCR in case of suspect colonies. Pseudomonas syringae pv. apii Agar method + PCR in case of suspect colonies. Pseudomonas syringae pv. coriandricola Agar method + PCR in case of suspect colonies. Dill, Coriander, Parsley - Detection of 2 pathogens Pseudomonas syringae pv. apii + Pseudomonas syringae pv. coriandricola Agar method + PCR in case of suspect colonies. Pseudomonas syringae pv. apii + Pseudomonas viridiflava Agar method + PCR in case of suspect colonies. | PA-BA-106 PA-BA-107 PA-BA-108 | 30 000 30 000 30 000 | 36 days 26 days | 278.00 289.00 359.00 |
| Pseudomonas viridiflava Agar method + PCR in case of suspect colonies. Pseudomonas syringae pv. apii Agar method + PCR in case of suspect colonies. Pseudomonas syringae pv. coriandricola Agar method + PCR in case of suspect colonies. Dill, Coriander, Parsley - Detection of 2 pathogens Pseudomonas syringae pv. apii + Pseudomonas syringae pv. coriandricola Agar method + PCR in case of suspect colonies. Pseudomonas syringae pv. apii + Pseudomonas viridiflava Agar method + PCR in case of suspect colonies. Pseudomonas syringae pv. coriandricola + Pseudomonas viridiflava | PA-BA-106 PA-BA-107 PA-BA-108 | 30 000 30 000 30 000 | 36 days 26 days | 278.00 289.00 359.00 368.00 |
| Pseudomonas viridiflava Agar method + PCR in case of suspect colonies. | PA-BA-107 PA-BA-108 PA-BA-109 | 30 000 30 000 30 000 | 36 days 26 days 26 days | |
| Pseudomonas viridiflava Agar method + PCR in case of suspect colonies. Pseudomonas syringae pv. apii Agar method + PCR in case of suspect colonies. Pseudomonas syringae pv. coriandricola Agar method + PCR in case of suspect colonies. Dill, Coriander, Parsley - Detection of 2 pathogens Pseudomonas syringae pv. apii + Pseudomonas syringae pv. coriandricola Agar method + PCR in case of suspect colonies. Pseudomonas syringae pv. apii + Pseudomonas viridiflava Agar method + PCR in case of suspect colonies. Pseudomonas syringae pv. coriandricola + Pseudomonas viridiflava Agar method + PCR in case of suspect colonies. | PA-BA-107 PA-BA-108 PA-BA-109 | 30 000 30 000 30 000 | 36 days 26 days 26 days | 278.0 289.0 359.0 368.0 |

Aromatic medicinal

| Pastoviology, Ungasted goods only | | _ | _ | _ |
|--|----------------------|--------------|-----------|---------|
| Bacteriology - Uncoated seeds only | | Size | Duration | Price |
| Dill, Coriander, Parsley | | 3.20 | Duration | |
| Candidatus liberibacter solanacearum | | | | |
| Detection by PCR. | PA-BA-CAND | 20 000 | 10 days | 136.0 |
| Mycology - See p.8 "Seed health" | | | | |
| Dill | | Size | Duration | Prid |
| Stemphylium botryosum, Alternaria radicina/carotiincultae (Stemphylium radicinu (Discolour section and other sections), Botrytis sp. | ım), Fusarium | | | |
| Agar method. | PA-ES-ANF | 400 | 19 days | 106.0 |
| Basil | | | | |
| Fusarium oxysporum , Fusarium (<i>Discolour</i> section and other sections), <i>Botrytis</i> sp. Agar method. | PA-ES-BAS | 400 | 19 days | 106.0 |
| Peronospora spp. | | | | |
| Grow-out test. | PA-MIBASGO | 400 | 42 days | 129.0 |
| Laurenden | PA-MIBASG3 | 3 000 | 42 days | 261. |
| Lavender | | | | |
| Phomopsis lavandulae, Botrytis sp., Fusarium (all sections), Phoma sp. Agar method. | PA-ES-LAV | 400 | 19 days | 106. |
| Parsley | ra-L3-Lav | 400 | 15 days | 100. |
| Septoria petroselini | | | | |
| Direct visual observation. UNTREATED seeds only. | PA-SE-PER | 1 000 | 15 days | 87. |
| Direct visual observation + counting. UNTREATED seeds only. | PA-SE-PERD | 1 000 | 15 days | 101. |
| Plasmopara nivea Seed wash method. UNTREATED seed only. | PA-MI-PER | 500 | 15 days | 102. |
| Alternaria petroselini (Stemphylium radicinum var. petroselini), Alternaria dauci, | | | | |
| Fusarium (all sections), Botrytis sp. | | | | |
| Agar method. | PA-ES-PER | 400 | 19 days | 106.0 |
| EVALUATION OF VARIETIES | | | | |
| Genotyping by molecular biology | | | _ | |
| | | Size | Duration | Pri |
| Рорру | | | | |
| /arietal identity control - SSR. | BI-G-BM-SSR-CID | | Contact E | BIOGEVE |
| Technological quality: biochemicals tests | | | | |
| Phovin | | Size | Duration | Pri |
| Stevia Steviosid and rebaudiosid A content by high performance liquid chromatography (HPLC). | BI-B-HPLC-STEV | | Contact E | BioGEVI |
| Field tests by SEV | | | | |
| | | | | Pri |
| DUS testing - Aromatic, Medicinal plants. | | SEV-DHS-ARON | NED | 2800.0 |

Micro-cleaning

icro cicariirig

Micro-cleaning of seed lots consists in determining the percentage of waste in raw seed lots, from a harvest, using sorting machines, laboratory replicates of industrial machines.

This activity enables the establishment of an optimal sorting diagram for the seed lot. It is an essential step in defining the industrial process for quality sorting in the factory, whatever the species. Moreover, the commercial value of a lot is estimated through precise knowledge of its quality.

HOW IT IS DONE?

Each species has his own morphological characteristics. Each morphological characteristic is associated with a sorting device, which settings are adjusted very precisely.

The complete sorting of a seed lot is carried out on a sorting line composed of several sorting machines ensuring complementarity on many criteria. In order to achieve the defined standards, the knowledge of characteristics, the expertise and the know-how of operators are essential.



Sorting on a raw batch of carrot before/after micro-cleaning

EQUIPMENTS

The SNES owns 20 different types of equipments in order to clean every types of seeds. Our training and expertise contribute to produce quality sorting, representative of the work provided in the factory. After the various sorting operations, analyses of specific purity and germination capacity can also be carried out at the SNES to ensure the quality of the seed lot.

| | | Price |
|--|----------|--------------|
| Micro-cleaning. Standard protocol. Beets . | MN-SN-01 | 59.00€ |
| Micro-cleaning. Standard protocol. Peas, Beans, Cucurbits. | MN-SN-02 | 54.00 € |
| Micro-cleaning. Standard protocol. Carrots. | MN-SN-03 | 84.00€ |
| Micro-cleaning. Standard protocol. Other vegetable crops. | MN-SN-04 | 76.00€ |
| Micro-cleaning. Standard protocol. Other field crop species. | MN-SN-05 | 67.00 € |
| Micro-cleaning. Standard protocol. Flower seeds. | MN-SN-06 | Contact SNES |
| Micro-cleaning. Mixed seeds. | MN-SN-07 | Contact SNES |
| Micro-cleaning. Quinoa. | MN-SN-08 | 100.00€ |
| Micro-cleaning. Standard protocol. Chicory | MN-SN-09 | 76.00 € |
| Micro-cleaning. Standard protocol. Small leguminous. | MN-SN-10 | 68.00€ |
| Additional charge for lots not presorted or requiring additional sorting time. | MN-SN-11 | 53.00€/h |
| Supplement fee. Details of each grid with percentage results. | MN-SUP | 12.60€ |

Requests for information: contact.mn@geves.fr






Evaluation of ploidy level from plants or seeds.


Cytology analyses carried out by the SNES aim to determine the level of ploidy by chromosome counting of root meristematic cells and/or flow cytometry. Ploidy defines the number of chromosome copies of a cell. The level of ploidy is characteristic of the species or variety. These analyses can be carried out from seeds or from plants on many species.

FLOW CYTOMETRY

Flow cytometry is a technic based on the marking of DNA with fluorochromes. The cytometer allows a precise measurement of the amount of fluorescence emitted by the cells after marking and excitation by a light beam. The measurement of the quantity of fluorescence emitted will then be compared to a control with a known level of ploidy. This will allow to conclude on the ploidy level of the tested sample.


Flow cytometry is mainly used to determine the level of ploidy of a series of plants and variety. In some cases, flow cytometer is also used to identify species with a very similar morphology or mutilated or poorly formed seeds.





MICROSCOPY

Chromosomal counting by microscopy is a technic that also makes it possible to define the level of ploidy. This is an essential step for species which do not have a reference for cytometry. Chromosome counting is carried out on meristematic root cells whose mitotic division has been blocked at the metaphase stage. The chromosomes are then observed and counted using a phase contrast microscope.



Metaphase cells of Festulolium



Metaphase cells of Gardenia

Requests for information or analyses: contact.cyto@geves.fr

Radiography 2D and tomography


Tools for evaluating seed quality.

WHY USE 2D OU 3D RADIOGRAPHY?


Radiography allows the internal morphology of seeds to be visualised. The objective is to understand or predict problems of physical or germinative quality. This tool also allows the phenotyping of precise characters of interest according to the request.

WHAT IS THE DIFFERENCE BETWEEN 2D RADIOGRAPHY AND TOMOGRAPHY?


2D radiography is a non-destructive method that allows rapid observation of different criterias on seeds (physical damages, empty seeds, insect damages, etc.). This technology allows a qualitative diagnosis of the state of the internal morphology. The Physical Analysis laboratory is ISTA accredited for these analyses.







Physical damages



Insect damages

3D radiography (tomography) is a technology whose method consists of generating a 3D image of the internal structure of an object. This tool applied to seeds allows the measurement of different characteristics and to obtain very precise quantitative data. The possible applications are diverse: characterisation of genotypes/varieties/batches, quantification of pathogen/insect damages, physical damages...



Evaluation of the quality of the coating



Quantification of insect damages



Quantification of cracks on a Corn seed

| | | Price |
|---|-----------|-------------------------|
| 2D radiography on seeds without interpretation (per digital image). | RX-IS-03 | 25.90 € |
| 2D image interpretation for internal morphological characterisation, the detection of insect/physical damage (%). | RX-SUP-03 | 16.20 € |
| Supply of one 2D image in .jpg format, for a particular determination or for measurements. | RX-SUP-RA | 1.10 € |
| For any request for information or analysis in 3D tomography: | RX-IS-05 | bea-tomographe@geves.fr |
| - Measurements of coating characteristics: | | |

- Measurements of coating characteristics
- Insect damages detection and associated volume measurements;
- $\hbox{-} \ \mbox{Measurement of internal seed constituents} \ ;$
- Measurement of seed filling rate;
- Detection and measurement of mechanical cracks and other damages;
- Other measures of interest.

| Visual or automatic image processing. | RX-SUP-05 | bea-tomographe@geves.fr |
|---|-----------|-------------------------|
| Supply of a batch of 2D images in jpg format. | RX-SUP-TO | bea-tomographe@geves.fr |

Biostimulation, Biocontrol, evaluation of treatment and the realization of tests under controlled conditions



GEVES, member of the Biocontrol Consortium and RMT BESTIM, provides its expertise for the characterization and evaluation of the effect of your treatments applied to seeds or seedlings.

Whether for biocontrol or biostimulant products, physical or chemical treatments, GEVES proposes to support you in the development of suitable evaluation methodologies and/or to carry out tests under controlled conditions. For *in vitro* and/or *in vivo* screening, or for the evaluation of disinfection, protection, stimulation or phytotoxicity effects, of treatment products in preventive and/or curative application.

SNES does not supply seeds or products. The sample size to be provided is 1 000 seeds per modality for selectivity and effectiveness assays. If only effectiveness trials are required, the sample size will be determined in relation to the project and the initial request.

GEVES is a multidisciplinary team of experts in seed quality and varietal resistance evaluation. It develops new evaluation methods in these areas that are recognized internationally. With this expertise, GEVES participates in research programs on biostimulation and biocontrol of seeds.

APPLICATION OF PRODUCTS ON SEEDS

Treatment of seeds is possible depending on the type of treatment and use. For more information, please contact SNES.

Depending on the quantity of seeds to be treated and the formulation of the product, 3 different tools can be used: Orbital agitator (20 g, liquid formulation); Hege bowl (500 g); Satec Concept treatment machine (up to 2 kg).

Application of a seed treatment product by SNES in the case of a treatment evaluation.

GE-APPLI

46.00

| SELECTIVITY TESTS | | |
|--|------------|-------|
| To check the selectivity of a treatment, the germination test should be determined on 400 seeds. | | Price |
| Vegetables. | GE-FG-18-4 | 65.00 |
| Cereals. | GE-FG-01-4 | 51.00 |
| Oilseeds. | GE-FG-17-4 | 54.00 |
| The percentage of seedlings showing phytotoxicity symptoms can be provided specifically. | | |
| All species. | GE-FG-PCPL | 23.10 |

EVALUATION OF TREATMENTS FOR SEED AND PLANT PROTECTION

| | | Contact |
|--|-------------|--------------------------|
| Evaluation of phytochemical products. | PA-EVAL-CHI | geoffrey.orgeur@geves.fr |
| Evaluation of biocontrol products, physical treatments and disinfection process. | PA-EVAL-BIO | |

| Few examples of available pathosystems ⁴ | | | | | | |
|---|---|-----------|---------------------------------|--|--|--|
| Wheat | Microdochium nivale | | Fusarium graminearum. | | | |
| | Tilletia caries. | Maize | Fusarium verticilioides. | | | |
| | Fusarium spp. (Fusarium graminearum, Fusarium | | Pythium ultimum. | | | |
| Villeat | avenaceum, Fusarium culmorum). | | Rhizoctonia solani | | | |
| | Puccinia striiformis, Puccinia triticina. | Sunflower | Botrytis cinerea | | | |
| | Pythium irregulare | | Plasmopara halstedii. | | | |
| | Plasmodiophora brassicae. | Sufflower | Verticillium dahliae. | | | |
| Rapeseed | Phoma lingam. | | Fusarium moniliforme. | | | |
| Kapeseed | Fusarium oxsporum conglutinans. | Cabbage | Hyaloperonospora brassicae. | | | |
| | Alternaria brassicicola. | Cabbage | Tryuloperonosporu brussicue. | | | |
| Beet | Aphanomyces cochlioide. | Lettuce | Fusarium oxysporum race 1 et 4. | | | |
| | Pythium sp. | Tomato | Meloidogyne incognita. | | | |

⁴Available pathosystems presented in evaluation of varieties as well as in seed health quality are all adaptable for evaluation of treatments.

EVALUATION OF BIOSTIMULANT PRODUCTS FOR GERMINATION AND/OR SEEDLING GROWTH

Two types of trials can be performed either under favourable conditions for the plant species (i.e. those applied in selectivity trials), or under penalizing conditions (i.e. abiotic stress).

| | Price / Contact | | |
|---|-----------------|---------------------------|--|
| Monitoring of seed germination on 200 seeds | | | |
| Germination energy (intermediate count; in addition to germination capacity). | GE-EG | 19.900 | |
| Counting dates for energy vary according to the species. | | | |
| Germination kinetics by image analysis (average rate of germination, kinetic curve). | GE-CI | sylvie.ducournau@geves.fr | |
| Seedling development tests | | | |
| Corn root length evaluation after 7 days germination at 15°C (4 replicates of 20 seeds). | GE-RAC | 77.00 | |
| Dry biomass of 4 replicates of 20 seedlings after germination test. | GE-BIOM | 55.00 | |
| Total length and root classification per diameter (4 replicates of 20 seedlings). | GE-CLASS | NEW 75.00 | |
| Growth kinetics by image analysis (Eloncam bench). | GE-ELON | sylvie.ducournau@geves.fr | |

Disease test supplies: inoculum and reference material

The available pests are listed on www.geves.fr. Specific preparation of isolate can also be done in the form of inoculum or artificially contaminated seeds. Warning: For the handling of quarantine pests, laboratories must be authorised to hold (Regulation 2019/829)

Pests' inoculum (contact SNES)

Tomato Rootstock.

| rests inoculant (contact SIALS) | | |
|---|------------|--------|
| | | Price |
| Specific preparation | | |
| Suspension of <i>Ditylenchus dipsaci larvae</i> (exemple of price: 1 335€ to inoculate 9000 plants). | PA-AD-DIT | / |
| Beet seedlings contaminated with viruliferous aphids <i>Myzus persicae</i> carrying yellowing virus BChV (<i>Beet chlorosis virus</i>). | PA-AD-MYZ | / |
| Other isolates and inoculum | | |
| One tray of 140 seedlings infected by a race of stripe/yellow rust (<i>Puccinia striiformis</i>). Contact jean-philippe.maigniel@geves.fr. | PA-AD-ROU2 | 130.00 |
| Inoculum supplied in Petri dishes. | PA-AD-INOC | |
| Inoculum supplied as contaminated cotyledons, plants or fresh leaves. | PA-AD-INOP | |
| Inoculum supplied in artificially contaminated grains that have lost germination capacity or artificially contaminated seeds that have maintained a germination capacity. | PA-AD-INOG | / |
| Inoculum supplied in liquid suspension. | PA-AD-INOL | 1 |
| Cyst of Globodera pallida ⁴⁰ or Globodera rostochiensis ⁴⁰ . | PA-AD-GLO | 1 |
| Cyst of Heterodera schachtii. | PA-AD-HET | |
| Reference material : isolates and seeds | | |
| | | Price |
| Bioagressors isolates | | |
| Specific preparation of reference isolate in Petri dishes (2 dishes/strain), dessicated (Bos) (1 g) or population of free living nematodes or cysts (around 20). | PA-AD-FOU | 168.00 |
| Specific preparation of 5 g of galls of <i>Meloidogyne incognita</i> (for inoculation of 15 to 20 plantlets). | PA-AD-MEL | 180.00 |
| Specific preparation of 5 g of galls of <i>Plasmodiophora brassicae</i> (for inoculation of 50 to 100 plantlets). | PA-AD-PLAD | 180.00 |
| Specific preparation | | |
| 100 mg of a vial of spores of stripe rust (<i>Puccinia striiformis</i>) or brown rust (<i>Puccinia recondita</i>) or crown rust (<i>Puccinia coronata</i>). | PA-AD-ROU | 62.00 |
| Specific preparation | | |
| 50 to 100 seeds of germinated Sunflower seeds contaminated by <i>Plasmopara halstedii</i> (downy mildew). | PA-AD-TOU2 | 180.00 |
| Lettuce seedlings infected with 1 race of <i>Bremia lactucae</i> , 30 cotyledons in the test period. | PA-AD-BREM | 180.00 |
| Erysiphe pisi, 2 seedlings with presence of sporulation. | PA-AD-ERYS | 180.00 |
| 2 cotyledons of Melon infected by 1 race of <i>Golovinomyces cichoracearum</i> (powdery mildew). | PA-AD-GOL | 180.00 |
| 2 cotyledons of Melon infected by 1 race of <i>Podosphaera xanthii</i> (powdery mildew). | PA-AD-POD | 180.00 |
| 2 Lettuce seedlings infected with <i>Nasonovia ribisnigri</i> race Nr: 0 with presence of apterae. | PA-AD-NAS | 180.00 |
| 30 leaves of Basil contaminated by <i>Peronospora belbahri</i> . | PA-AD-BEL | 180.00 |
| Controls/differential hosts vegetables (MATREF) for one sowing unit (1 g for Bremia, 200 seeds for other pathogens) | | |
| Complete pack of differential hosts for <i>Bremia</i> of Lettuce . | PA-HD-BLAI | 366.00 |
| Carrot. | PA-HD-CAR | 50.00 |
| Squash. | PA-HD-COU | 88.00 |
| Watermelon. | PA-HD-PAS | 88.00 |
| Bean. | PA-HD-HAR | 69.00 |
| Lettuce. | PA-HD-LAI | 69.00 |
| Corn salad. | PA-HD-MAC | 50.00 |
| Melon. | PA-HD-MEL | 88.00 |
| Capsicum. | PA-HD-PIM | 101.00 |
| Pea. | PA-HD-POI | 69.00 |
| Tomato. | PA-HD-TOM | 88.00 |
| | | |

PA-HD-PGTO

101.00

Sector support



Inter-laboratory proficiency testing (ILPT) is used to evaluate the ability of a laboratory to perform a method. For more information, visit our website www.geves.fr.

The organisation of comparative tests includes planning and delivery of documents to participants, preparation of samples, definition of a reference, interpretation of results and issuing of a final report.

Not included: supply of seeds cost (billed at actual price).

Inter-laboratory proficiency tests – PT & Other comparisons

| | Price | Contact |
|---|--------------|-----------------------|
| Purity – All species (based on 15 participants) – by sample. | 186.00 | |
| Germination – All species (based on 15 participants) – by sample. | 128.00 | |
| Moisture content – All species (based on 15 participants) – by sample. | | |
| Thousand-seed weight – All species (based on 15 participants) – by sample. | | Fabienne BRUN |
| Seed health. | Contact SNES | eil.semences@geves.fr |
| Organisation of inter-laboratory comparisons tests on request. | Contact SNES | |
| Supply of reference samples for internal laboratory control. | Contact SNES | |
| Expertise in the case of atypic results on seeds assay or deviation found (control card for recognized laboratories). | Contact SNES | |

AUDITS

According to various standards (ISTA, recognition in the context of certification), laboratory audits can be carried out to analyse your organisation. One-day audit includes an analysis of a pre-audit file, the conducting of the audit as well as the audit report.

Contact: Fabienne BRUN (audit.semences@geves.fr).

REFERENCE MATERIALS AND DOCUMENTS SUPPLIES

Find all our publications and reference materials in the different chapters of the price list and on our website www.geves.fr.

TRAININGS - EXPERTISES

| To apply for training | | Price | Contact |
|--|---------|--------------|--------------------------------|
| Technichal training with SNES. | | Contact SNES | Fabienne BRUN |
| Seed quality analysis, inter or in-company, at SNES or on-site. | | | formation.semences@geves.fr |
| Technichal training with BioGEVES. | | Contact SNES | biogeves.analyses@geves.fr |
| Technichal training with SEV. | | Contact SNES | rachel.tessier@geves.fr |
| For the setting up of an expertise in an international context | | | |
| Technical expertise and visit. | | Contact SNES | secretariat.direction@geves.fr |
| Collective reading of results | | | |
| Collective reading of germination results, details of abnormals and debriefing of the results reading, per sample. | GE-LECT | 102.00 | Inr.semences@geves.fr |

Terms and Conditions



Article 1 - General Information

The present general terms and conditions of sale apply for services which appear in the GEVES price list (Variety and Seed Study and Control Group), public interest group governed by the constitutive convention of July 17, 1989, having made the object of an approval order dated July 17, 1989 and its modified constitutive convention of April 17, 2014 whose head office is located 25 rue George Morel, CS 90024, 49071 Beaucouzé Cedex FRANCE.

The main official missions of GEVES are to conduct studies or analyses of:

- characterization and/or identification of varieties,
- agronomic quality of varieties,
- physical, physiological and sanitary control of seed.

Article 2 - Object and field of application

The analyses carried out within the framework of any order are in accordance with the present general terms of sale.

The placing of an order implies full acceptance of these general terms of sale which prevail on any other document of the customer, unless otherwise agreed between the customer

Geves reserves itself the right to modify the present general terms of sale.

Article 3 - Orders

3-1) Order taking

The orders are definitive only when the present general terms of sale are full accepted by the legal representative of the customer or any person duly appointed for that purpose

The customer has to respect the terms of the supply of material described in the GEVES price list.

3-2) Modification of the order

The terms of the orders transmitted to GEVES are irrevocable for the customer, except written acceptance from GEVES. On this assumption, GEVES will not be held anymore by the deadlines agreed upon at the moment of the initial order.

If a customer places an order to GEVES, without having carried out the payment of preceding orders despite reminder from GEVES, GEVES can repudiate the order, without the customer being able to claim any allowance, whatever the reason.

GEVES reserves itself the right to refuse any order.

Article 4 - Delivery of the results

4-1) Delivery time

The delivery time of the results are given only on a purely informative and indicative basis; those depending in particular on arrival of the orders, the respect of the conditions of preparation of the samples sent by the customer (weight, number, packing for example), request for more information, or complementary analyses. For each service, useful information is available on the GEVES website (www.geves.fr). In any assumption, the delivery within the deadlines can intervene only if the customer is up to date of his obligations with GEVES.

GEVES shall endeavor to meet agreed deadlines with the customer.

Delays of delivery of results cannot lead to any penalty or allowance, nor to justify the cancellation of the order.

4-2) Terms

The delivery of the results is made by paper form or by electronic way.

4-3) Complaints

The complaints are to be forwarded to the customer service of GEVES whose contacts appear in the GEVES price list. GEVES acknowledges to the customer the receipt of the complaint, deals with it and defines an appropriate treatment as soon as possible. GEVES shall inform the plaintiff of the progress of the claim and the conclusions.

Except explicit indication of the customer validated by the customer service of GEVES whose references are indicated on the GEVES price list, no material submitted for analysis will be

Article 6 - Guarantee - Liabilities

6-1) Scope

GEVES provides services. As such, GEVES is under the obligation of best effort. It could not be held responsible for non-satisfactory results from the point of view of the customer, for causes of which it does not have the control. GEVES will have, if necessary, to issue reserves

6-2) Exclusions

If the elements provided by the customer do not allow the fulfillment of the ordered service, GEVES will inform the customer. If this situation persists, the liability of GEVES could in no way be required.

In particular, GEVES could not be held responsible for sampling (except for Orange ISTA Certificates for which GEVES is responsible for sampling), the collecting, the conditioning and the transport of the samples, which is the customer's entire liability. Moreover, the samples received at GEVES shall be in good condition of conservation and shall not present identified risk for the staff of GEVES or for the environment. When a phytosanitary treatment has been applied, the customer shall inform GEVES

The customer waives all right to take any action against GEVES for all losses or all direct or indirect damages resulting from the services, as well as in the situation where the services of GEVES would be unsuitable for the uses of the customer.

The rates applied to the orders are those indicated in the GEVES price list, unless particular conditions negotiated with GEVES.

Any order made on the basis of a quotation established by GEVES will be taken into account only after signature of the quotation, by the legal representative of the customer or any

person duly elected for that purpose.

Prices are indicated exclusive of VAT, based on current rates and will be increased by current taxes of all types on the invoicing date.

Amounts are indicated in Euros. Payments should be made in Euros.

The transport fees of the samples provided to GEVES for analysis are always at the charge of

Article 8 - Invoicing

Any order, even if it is cancelled during the execution of the service, will give rise to an invoice. Elements of identification of the customer and ordered services are indicated on the invoices. The customer service of GEVES whose references appear in GEVES price list can be contacted for any question related to the invoice.

Article 9 - Payment

9.1) - Time for payment

The maximum payment time is 60 days from the date of emission of the invoice.

9.2) - Terms

The payments shall be made:

- by French postal or bank check or credit or postal transfer addressed to: GEVES, 25 rue George Morel, CS 90024, 49071 Beaucouzé Cedex FRANCE

- by signed and accepted draft or promissory note. GEVES does not authorize any discount for cash payment or on a former date to those resulting from these general terms of sale.

9.3) - Delay of payment

Any sum still not paid at the due date by the customer will give rise to the payment of penalties at the rate of the European Central Bank plus 10 points and a lump sum of 40 Euros for recovery costs in compliance with Decree n° 2012-1115. These penalties are payable automatically without prior notice from GEVES on the date following the due date. Moreover, GEVES reserves itself the faculty to apply to the competent court of law to stop this non-fulfillment, under penalty per day of delay.

Article 10 - Confidentiality - Rights of ownership

GEVES guarantees the confidentiality of the results of analysis, unless the detection of a quarantine pathogen. Under such circumstances, GEVES has to communicate immediately to the qualified services of the ministry in charge of agriculture all information relating to the material in which the quarantine pathogen was identified.

This exception also applies to other situations, such as the detection of fortuitous presence of GMO, if the regulation in force imposes to GEVES to communicate information to the qualified services of the French State.

The results provided by GEVES can in no way being modified, reproduced or diffused even in a partial way, to third party, without the preliminary authorization of GEVES. Duplicates can be obtained on request at the customer service of GEVES whose references are indicated on GEVES price list.

Article 11 - Personal data

For any processing of personal data carried out in connection with this Quotation, the Parties shall comply with Regulation (EU) 2016/679 of the European Parliament and of the Council of 27 April 2016 on the protection of individuals with regard to the processing of personal data and on the free movement of such data, as transposed into French Law No 2018-493 of 20 June 2018.

Each Party represents and warrants to the other Party that it will strictly comply with GDPR for any processing of personal data in connection with this Quotation.

Personal data collected and processed by the Parties in the context of this contractual relation are necessary for its execution (legal basis). They are kept for a period of 10 years (retention period) from the date of the end of the Quotation.

Article 12 - Agreement of proof

In accordance with Articles 1316-1 to 1316-4 of the Civil code, documents in electronic form are admitted as evidence in the same way as paper-based documents

The Parties expressly agree that this Quotation concluded in electronic form and signed in a dematerialized way, as well as the documents relating to it:

- Constitute the original documents:
- Are drawn up and kept under conditions that guarantee their integrity;
- Are perfectly valid between them. As such, the Parties undertake not to challenge the validity, enforceability or probative value of this Quotation and the documents relating to it on the basis of their conclusion or transmission by electronic means;
- Constitute written evidence within the meaning of the aforementioned Articles 1316-1 to 1316-4 of the Civil Code. Thus, this Quotation concluded by electronic means is deemed to be evidence of the content of the Quotation, of the identity of the signatories and of their consent to the obligations arising from the Quotation.

Article 13 - Force majeure

The emergence of a case of force majeure causes the suspension of the execution of the

Article 14 - Attribution of jurisdiction

For all disputes relating to the services carried out by GEVES, including those relatives to the interpretation of the general terms of sale, the jurisdictions of Angers shall be qualified.

Article 15 - Applicable law

The present general terms of sale, and any question which it would omit to treat, shall be exclusively governed by the French law.

By appending his signature on the Quotation, the customer:

- recognizes and accepts without reserve the present general terms of sale and that those will apply to all the further orders until communication of new general terms of sale by
- declares that he has read and accepts them,
- waives its own purchasing conditions

OUR PUBLICATIONS

AND REFERENCE MATERIAL







More information at www.geves.fr

Contact: Inr.semences@geves.fr



Groupe d'Étude et de contrôle des Variétés Et des Semences